

## **REFERENCES**

## REFERENCES

- Abir, S.(2014).Seasonal variations in physico-chemical characteristics of Rudrasagar wetland- A Ramsar site, Tripura, North East, India. *Research Journal chemical Sciences*, 4(1):31-41
- Abujam S.S.; Shah R.K.; Deori, D. J. and Biswas, S.P. (2014). Diversity of Macrophytes in Maijan Wetland, Upper Assam, India. *South Asian Journal of Experimental Biology*, 4(4): 164-171.
- Achikanu, C.E.; Eze-Steven, P.E.; Ude, C.M. and Uguriokolie O.C. (2013). Determination of the vitamin and mineral composition of common leafy vegetables in south eastern Nigeria. *International Journal of Current Microbiological and Applied Sciences*, 2(11): 347-353.
- Adelakun, K.M.; Kehinde, A.S.; Amali,R.P.; Ogundiwin, D.I.andOmotayo, L. (2016).Nutritional and Phytochemical Quality of some Tropical Aquatic Plants. *Poultry, Fisheries and Wildlife Sciences*, 4(2):2-4.
- Adeyeye, E.I. and Otokiti, M.K.O. (1999).Proximate composition and some nutritionally valuable minerals of two varieties of *Capsicum annum* (Bell and Cherry Peppers). *Discovery and Innovation*, 11(1): 75- 81.
- Adikari, A.M.A.N.; Sundarabarathy, T.V.; Herath, HMUKPB; Nayananjali, W.A.D and Adikari, A.M.J.B. (2017). Formulation of artificial feeds for Indian Carp (*Catla catla*) fry using aquatic plants (*Ipomea aquatic* and *Hydrilla verticillata*). *International Journal of Scientific and Research Publications*, 7(7): 83-89.
- Adoni, A.D. (1975). Studies on microbiology of Sagar Lake. Ph. D. Thesis. H.S. Gour University, Sagar. M.P., India, 1-254 pp.
- Agarwala, N.K. (1996).Limnology and fish Productivity of Tamranga wetland in Bongaigaon District of Assam (India) with special reference to some productivity indicator. Ph.D. Thesis, Gauhati University.

- Ahirrao, S.D. and Mane, A.S. (2000). The diversity of Ichthyofauna, taxonomy and fisheries from freshwater of Parbhani, Dist. Maharashtra State. *Journal Aquatic Biology*, 15(1&2): 40-43.
- Ahmed, K.K.; Haque, M.K.I. and Haque, M.E. (1999). Studies on some Physico-chemical factors of Kaptai Lake. *Bangladesh Journal of Fisheries Research*, 3(1): 33-39.
- Akram, Z.; Fatima, M.; Shah, S.Z.H.; Afzal, M.; Hussain, S.M.; Hussain, M. (2019). Dietary zinc requirement of *Labeo rohita* juveniles fed practical diets. *Journal of Applied Research*, 47(1):223-229
- Allen, M.B. (1966). Distribution of the chlorophylls. In L.P. Vernon and G.R. Seely eds., *the chlorophylls*, New York: Academic Press.
- Al-Thagafi, Z.; Arida, H.; Hassan, R. (2014). Trace toxic metal levels in canned and fresh food: a Comparative study. *International Journal Innovative Research in Scence, Engineering and Technology*, 4(3): 8977-89.
- Alikunhi, K.H. (1957). “Fish Culture in India” FM. Bull. No. 20, ICAR, New Delhi.
- Ambasht, R.S. (1990). A Text book of plant ecology, Tenth edition, Students Friends & Co, Varanasi, 373 pp.
- Ambasht, R.S. (2005). Macrophytes limnology in the Indian subcontinent, Ukaaz Publication, Hyderabad, 58-174 pp.
- Anderson, R.A. (1981). Nutritional role of Chromium. *Science of the Total Environment*, 17(1):13- 29
- AOAC (2005). Official methods of Analysis, 17<sup>th</sup> edition, Association of the Analytical Chemists, inc, Virginia, USA, 1-40 pp.
- Anon, (1967). Making aquatic weeds useful: Some perspectives for developing countries. National Academy of Sciences. Washington, D.C., 1-175 pp.

APHA (2005).Standard method for the Examination of water and waste water, 21<sup>st</sup> edition, American Public Health Association,Washington DC, USA

Arivozhagan, P.and Kamalaveni K. (1997).“Seasonal variation in physic-chemical parameters and plankton analysis of Kurichi pond”. *Journal Environment and Ecology*, 15(2): 272-274.

Arun, L.K. (1997). *Patterns and processes of fish assemblage in Periyar lake valley system of Southern western Ghats, KFRI Research Report.* Peechi, Kerala, Kerala Forest Research institute

Arnon, D.I. (1949). Copper enzymes in isolated chloroplasts, Polyphenoloxidase in *Beta vulgaris*. *Plant Physiology*, 24(1): 1-15.

Bag, M.P.; Mahapatra, S.C.; Rao, P.S. and Chakrabarty, D. (2011). Making aquatic weed as potential feed for Nile tilapia (*Oreochromis niloticus* L.) and its impact on fatty acid profile. *International Research of Pharmacy and Pharmacology*, 1(8):194-202.

Bag, M.P.; Mahapatra, S.C.and Rao, P.S. (2012). Aquatic weed as potential Feed for Mozambique tilapia, *Oreochromis mossambicus*. *Journal of Aquaculture Research & Development*, 3(8): 1-4.

Banarjee, S.M. (1967). Water quality and soil conditions of fish ponds in some states of India in relation to fish production. *Indian Journal of Fisheries*, 14: 115- 125.

Banerjee, G.C. (1988). Feeds and principles of Animal nutrition(2<sup>nd</sup>. Edn.) A rapid method of total lipid extraction and purification. *Canadian journal of Biochemistry and physiology*, 37: 911-917

Banerjee, A.and Matai, S. (1990). Composition of Indian aquatic plants in relation to utilization as animal forage. *Journal of Aquatic plant Management*, 28: 69-73

Barooah, C.and Sarma, L.(2016).Vertebrates of Assam- A checklist with IUCN status, Published by Assam Science Technology and Environment Council, Bigyan Bhawan, G.S. Road, Guwahati, Assam, India, 14-313 pp.

- Barbuiya, A.H.; Bhattacharyya, R.C. and Dutta, A. (2015). A Study of the Limnology and Ichthyology of Dhir Beel at Dhubri, Assam, India. *International Research Journal of Biological Sciences*, 4(8):40-48.
- Barbaruah, A.D.; Paul, S.; Kalita, B. and Dutta, A. (2012). Studies on Fish Diversity, Plankton and Macrophytes of Monoha Beel in Morigaon, Assam. Published by Dr. S.K. Dutta, Principal/ President, National Seminar Organizing committee, Dhing College, Nagaon, Assam. 158-165 pp.
- Bardach J.E.; Ryther, J.H. and Melarney, W.O. (1972). Aquaculture: The farming and husbandry of freshwater and marine organisms, Wiley-Interscience, New York
- Barion, G.P.; Blane, B.; Chassing, P.; Olive and Pelletier, G.R. (1982). "Quality of Water in Lake Lemen. *France Eakque*, 15(2):154-157.
- Baruah, P.; Gogoi, R.; Bordoloi, S.; Ahmed, S.; Sharma, K.; Goswami, D.C.; Murthy, TVR.; Shing, T.S. and Garg, J.K. (1997). Wetland of Assam: A Project report; Assam Remote Sensing Application Centre, Guwahati and Space Application Centre (ISRO), Ahmedabad
- Baruah, B. K. and Barthakur, C.R. (1998). "Water quality of ponds in Chandrapur area of Kamrup district, Assam". *Journal of Environment and Ecology*, 16(2): 254-256
- Baruah, P.P. (2003). Phytosociological account of macrophytes of a disturbed Lentic Habitat in middle Assam. *Annals of Biological Research*, 11(1): 27-36.
- Baruah, D.; Borah, S.; Hazarika, L.P.; Dutta, R.; Bakalial, B.; Biswas, S.P. and Sarma S.K. (2011). A simple diagnostic tool for measuring river health-example from a tropical snow fed river. *Annals of Biological Research*, 2(5): 432-443.
- Barman, D.; Roy, B.; and Roy, S. (2015). Seasonal variation of physico-chemical characteristics of Wetlands in the West Garo Hill, Meghalaya, India. *International Research Journal of Biological Science*, 4(1): 60-65.

- Bates, B.C.; Kundzewicz, Z.W.; Wu, S.; Palutikof, J.P.(Eds.)(2008). Climate Change and Water [Technical paper VI]. Intergovernmental panel on climate change, Geneva.
- Beaver, R.J.; Thomas, L.; Cristoman, L.and James, S. Bays (1981).“Thermal regimes of Florids (USA) lakes.” *Hydrobiologia*, 83(2): 267-274.
- Beavan, R. (1877).“Handbook of freshwater fishes of India.” London: L. Reeve &Co (reprinted 1972, Narendra Publishing House, Delhi, 247 pp., 8 Pls.
- Berra, T.M. (2001). Freshwater fish distribution. Academic Press, San Diego, C.A
- Best EPH (1982). Effect of water pollution on freshwater submerged macrophytes. Water pollution and Management review, 27-56 pp.
- Bhagabati, A.K.;Kalita,M.C.and Baruah,S.(2006). Biodiversity of Assam, Status, strategy and Action plan for conservation. Published by EBH Publishers (India), 136, M.L.N., Road, Panbazar, Guwahati, Assam (India), 107-111 pp.
- Bhagabati, A.K.; Kar, B.K. and Bora A.K. (2011). Geography of Assam. Published by Rajesh Publications, 1.Anarsi Road, Darya Gang, New Delhi, 1-315 pp.
- .Bhakta, J.N. and Bandyopadhyay, P.K.(2008). Fish Diversity in Freshwater Perennial water bodies in east Midnapore district of West Bengal, India. *International Journal of Environmental Research*, 2(3): 255-260.
- Bhaskar, V.and Raji, B.A. (1973). Hydrophytes and marsh plants of Mysore city. Prasaranga, University of Mysore, Mysor, India
- Bhattacharjya, B.K.; Sugunan, V.V. and Choudhury, M. (2000). Threatened fishes of Assam, Pp. 75-79. In: Ponniah, A.G. and Sarkar, U.K. (eds). Fish Biodiversity of Northeast India, NBFGR, Lucknow, NATP Publ. 2-228 pp.

- Bhattacharjya, B.K.; Choudhury, M. and Sugunan, V.V. (2003). Ichthyofaunistic resources of Assam with a note on their sustainable utilization, pp.87-105. In: Participatory Approach for fish biodiversity conservation in North East India. Pub. Director, NBFGR, Lucknow, India.
- Bhattacharjee, D.K.; Sarma, S.K., Devi, B. (2001). Medicinal plants used by Bodo tribe of Nalbari district in Assam. *Ethnobotany*, 13: 135-139.
- Bhattacharjee, D.K.; Sarma, S.K., Devi, B. (2002). Traditional use of herbal medicines by Madahi tribe of Nalbari district of Assam. *Ethnobotany*, 14(1&2): 103-111.
- Bhattacharjee, D.K.; Sarma, S.K., Devi, B. (2006). Ethnomedicinal uses of plants by the Sarania tribe of Nalbari district of Assam. *Journal of Economic and Taxonomic Botany*, 30(Suppl.): 133-139
- Bhattacharjee, D.K.; Sarma, S.K.; Bora, P.C. and Kar, A. (2008). Notes on selected herbal treatment of the common people in Barpeta district, Assam. *Journal of Advance Plant Sciences*, 4 (1&2): 69-73.
- Bhattacharjya, D.K. and Borah, P.C. (2008). Medicinal weeds of crop fields and role of women in rural health and hygiene in Nalbari district, Assam. *Indian Journal of Traditional Knowledge*, 7 (3), 504- 501.
- Bhaumik U.; Das, P. and Paria, T. (2004). Macrophytes and Macrophyte Associated Fauna in a flood plain wetland system of West Bengal having both open and close characteristics. *Environment and Ecology*, 22(Spl.-2): 371
- Bhosale, S.V.; Bhilave, M.P. and Nadaf, S.B. (2010). Formulation of Fish Feed using ingredients from plant sources. *Research Journal of Agricultural Sciences*, 1(3): 284-287.
- Bhowmik, M.L. and Ayyappan, S. (2000). Biodiversity Conservation in North east India-a Challenge. P.71-72. in: A.G. Ponniah and U.K. Sarkar (eds.) Fish biodiversity of North east India. NBFGR. NATP Publ. 2228 pp.

- Bhowmik, S.; Saha, R.and Datta, B.K. (2008). Aquatic and marshland plants in West Tripura, India, *Pleione*, 2(1): 3-11.
- Bhowmik, S.; Dutta, B.K.and Saha, A.K. (2012). Determination of mineral content and heavy metal content of some traditionally important aquatic plants of Tripura, India using atomic absorption Spectroscopy. *Journal of Agricultural Technology*, 8(4): 1467-1476.
- Bhute, K.B. and Harney, N.V.(2017). Macrophytes biodiversity of Nagrala lake of Bhadrawati, district-Chandrapur (M.S.), India. *International Research Journal of Science & Engineering*, 6(1):17-19.
- Bhuyan, K.C.; Dutt, A.and Kalita, B.(2009). Hydrobiology and Fishery Status of Sondoba Beel, Morigaon, Assam. *Journal of Inland Fisheries Society of India*, 41(2): 48-53.
- Billiore, D.K.and Vyas, L.N. (1981).Distribution and production of macrophytes in pichhola Lake, Udaipur (India). *International Journal of Ecology and Environmental Sciences*, 7:45-54
- BIS (1993)."Indian Standards for surface water, Bureau of Indian Standards" New Delhi, IS 2296.
- Biswas, K.and Calder, C. (1937). Handbook of common water and marsh plants of India and Burma. Bishen Singh & Mahendra Pal Singh, Dehradun
- Biswas, S. P.and Choudhury, M. (2008). Ecology and Ichthyofaunal Diversity of wetlands in upper Assam. In: Management of Fresh water Ecosystem, Agrotech Publishing Academy, Udaipur.73-82 pp.
- Biswas, S.P. and Boruah, S.(2002). Problems and prospects for the development of fisheries in the Brahmaputra basin. In: Fish germplasm inventory and conservation in North East India (Eds. A. G. Ponniah and U.K. Sarkar), NBFGR, Lucknow.
- Biswas, K. and Calder, C.C. (1984). Hand book of common water and marsh plants of India and Burma. 2<sup>nd</sup> ed. Bishen Singh and Mahendra Pal Singh, Dehradun, 1-216 pp.

- Bone, Q. and Marshall, N.B. (1983).The Mauthner system of deep sea teleosts. *Journal of the Marine Biological Association of the United Kingdom*, 63(4): 881-896.
- Bora, S. and Biswas, S.P.(2015). Water quality and ichthyofaunal diversity of an oxbow lake in upper Assam. *International Journal of Fisheries and Aquatic Studies*, 3(1): 15-18.
- Borah J., Choudhury E. and Choudhury N.K. (2017). A Preliminary Study on the Ichthyofaunal Diversity of Kankati Beel of Biswanath District, Assam. *IJSRD-International Journal for Scientific Research & Development*, 5(06): 50-53.
- Bordoloi, R. (2012). Comparative study on fish and fisheries of a closed and an open type wetland of the Brahmaputra valley. *International Journal of Science and Research Publication*, 4(2):1-11
- Bordoloi, R.and Hazarika, A.K. (2015). Biodiversity and Conservation Status of Ichthyofauna of Doria Beel, Majuli, India. *Research Journal of Animal, Veterinary and Fishery Sciences*, 3(8): 1-8.
- Borker, M.R.; Saraswat, K.; Quadros, S.V.and Fernandes, D.(1992).“Diel cycle of some abiotic and microbial qualities of a freshwater reservoir during monsoon at Marmugoa, Goa”. *Bioved*, 3(2): 133-136.
- Borthakur, S.and Sharma, K. (1998). Effect of some non- conventional fish meal replaces on the growth feed conversion and body composition on *Clarias batrachus* (Linn.) fingerlings. *Environmental Ecology*, 16(3): 694-698.
- Boulton, A.J.and Brock, M.A. (1999). Australian freshwater ecology: Processes and management, Gleneagles publishing, Adelaide.
- Boyd, C.E. (1968). “Fresh water plants: A potential source of protein”. *Economic Botany*, 22: 359-368.
- Boyd, C. E. (1969).“The nutritive value of three species of water weeds”. *Economic Botany*, 23: 123-127.

- Boyd, C.E. (1970).Vascular Aquatic plants for Mineral nutrient removal from polluted waters. *Economic Botany*, Vol.xxIII (1), 95-103 pp.
- Boyd, C.E. (1982).Water quality Management for Pond Fish Culture. Developments in Aquaculture and Fish Science. New York, U.S.A.: Elsevier Sci. Publ. Co.
- Boyd, C.E. (1984).Water quality Management in Aquaculture. Special Publ.No.22, Cent.Mar.Fish.Res.Inst., Kochi,India.
- Boyer, R. (1990). Isolation and Spectrophotometric Characterization of Photosynthetic Pigments. *Biochemical education*, 18(4): 203-206.
- Bowen, W.R. and Baxter,W.D.(1980). Experimental Cell Biology, An Integrated Laboratory Guide and Text, Macmillan Publishing Co, Inc., New York.
- Brahma,N. K. and Misra,A.K. (2012). Hydro-Geochemical study of Floodplain wetland: A case study with Respect of the seasonal variation of Rupahi Beel, Nagaon, Assam. Published by Dr. S.K. Dutta, Principal / President, National Seminar organizing Committee, Dhing College, Nagaon, Assam. 213-226 pp.
- Briggs, J.C.(2005).The biogeography of Otophysan fishes(Ostariophysi: Otophysi); A new appraisal. *Journal of Biogeography*, 32: 287-294.
- Britton, G. (1983).The biochemistry of natural pigments.Cambridge University Press. 133-140 pp.
- Brody, T. (1994).Nutritional Biochemistry. San Diego, Academic Press, 1-1006 pp.
- Brown, S.B.; Houghton, J.D.and Hendry, G.A.F. (1991). Chlorophyll breakdown. In Scheer H (Ed.): Chlorophylls. Boca Raton, CRC Press, 465-489 pp.
- Burlakoti, C. and Karmacharya, S.B.(2004).Quantitative investigation of aquatic macrophytes in Beeshazar Tal (Beeshazar lake). *Himalayan Journal of Sciences*, 2(3): 37-41.

- Butnariu, M. (2016)."Methods of Analysis (Extraction, Separation, Identification and Quantification) of Carotenoids from Natural products". *Journal of Ecosystem & Ecography*, 6(2): 1000193`
- Calvin, M. (1959). From microstructure of macrostructure and function in the photochemical apparatus. In The photochemical Apparatus- its structure and function. *Brookhaven Symposium Biology*, 11:160.
- Chacko, P.I. (1948). Development of fisheries of Periyar Lake. *Journal Bombay Natural History Society*, 48(1): 191-192.
- Chakraborty, S.; Kausor, M.A.; Goyal, A. K. and Basumatary, A.K. (2017). Studies on Physico-Chemical Parameters of Fish Inhabiting Four Water Bodies in the Vicinity of Kokrajhar, BTAD, Assam. *International Journal of Applied Chemistry*, 13(3): 409-420.
- Chakrapani, B.K.; Ramakrishna Parma, V. R. (1996). Quality of lake waters in and around Bangalore and Maddurai. Water birds and wetlands of Bangalore
- Chakraborty, P. (2018). Review in freshwater fish diversity of Northern West-Bengal India: Status, threats and conservation measures. *Intnernational Research Journal of Biological Science*. 7(7): 39-51.
- Chakravarty, P.; Chakravarty, M.and Sharma,S. (2012). A Survey on the Fish Diversity with Special Reference to the Classified Ornamental Fishes and their Prospects in the Kapla Beel of Barpeta District. *The Science probe.*, 1(2): 12-21.
- Chandra, S.; Singh, A. and Tomar, P. K. (2012). "Assessment of water quality values in Porur Lake Chennai, Hussain Sagar Hyderabad and Vihar Lake Mumbai, India". *Chemical Science Transactions*, 1(3): 508-515.
- Chavan, A.R. and Sabnis, S.D. (1961). A study of hydrophytes of Baroda and environs. *Journal of the Indian Botanical Society*, 40(1): 121-130.

- Chen, B.H. and Chen, Y.Y. (1992). Determination of Carotenoids and Chlorophylls in water convolvulus (*Ipomoea aquatic*) by liquid Chromatography. *Food Chemistry*, 45:129-134.
- Chen C.Y., Stemberger R.S., Klane B., Blum J.D., Pickhardt P.C. and Foll C.L. (2000). Accumulation of heavy metals in food web component across a Gradient of Lakes. *Limnology and Oceanography*, 45(7): 1525-1536
- Chesters, J.K. (1991). Trace Elements-Gene Interaction with Particular Reference to Zinc. *Proceeding of the Nutrition Society*, 50(2): 123-129 pp.
- Chhetry, B. and Deka, P.(2016). Ichthyofaunal Diversity of era Kopili Beel of Karbi angling District of Assam, India. *International Journal of Advanced Science and Research*. 1 (6): 33-41.
- Chilgar, O.S. and Jagtap,H.S.(2018).Study of Ichthyofaunal diversity of Marathwada region,Maharashtra with references to *Clarias batrachus*. *Trends in Fisheries Research*, 7(2):81-86.
- Chitranshi, V. R. and Bilgrami, P. A. (1986). "Comparative ecological studies on two ox-bow lakes of river Burhi Gandak. 1. Report on macrophytes". *Proceedings of the National Academy of Sciences, India Section B*, 56(3): 247-253.
- Cho, C. Y. (1985). Effects of protein intake on metabolically and net energy value of fish diet Pp. in nutrition and feeding in fish, C.B. Cowry, A.M.Mackie and J.G. Bell, Eds. London: Academic Press. 95-117 pp.
- Chovanec, A.; Hofer,R. and Schiemer,F.(2003).Bioindicators and Biomonitor. Elsevier Publisher, 5215(3): 80148-0.
- Chowdhury, M. (2009).Plant Diversity and Vegetation Structure in the Wetlands of Malda District of West Bengal, India. Ph.D. Thesis, North Bengal University.
- Chowdhury, M. and Das, A.P. (2010). Hydrophytes of different wetlands in the Maldah District of West Bengal, India. *Environmental Biology Conservation*, 15: 22-28.

- Chowdhury, M. and Das, A.P. (2011). Macrophytic diversity and community structure of *Adh Soi* wetland of Maldah district of West Bengal, India. In C. Ghosh& A.P. Das, *Recent Studies in Biodiversity and Traditional Knowledge in India*. Sarat Book House, Kolkata. 109-115 pp.
- Choudhary, J.; Singh, S.N. and Singh, S. (2014). Physico-chemical and Biological Parameters of the Three Rural Ponds of Sasaram of Bihar. *Int. Journal of Applied Sciences and Biotechnology (IJASBT)*, 2(1): 206-210.
- Clark, J.; MacDonald, N.L. and Stark, J.R. (1987). Leucine aminopeptidase in the digestive tract of Dover sole [*Solea solea* (L.)]. *Aquaculture*, 61(3-4): 231- 239.
- Cook, C.D.K. (1996). Aquatic and Wetland Plants of India. Oxford University Press, NewYork, 385 pp.
- Cunningham, L.W.; Fischer, R.L. and Vestling C.S. (1955). A Study of the Binding of Zinc and Cobalt by Insulin. *Journal of the American Chemical Society*, 77(21): 5703- 5707 pp.
- Conard, H.S. (1905). The water lilies: A monograph of the genus *Nymphaea carnegiei* Inst. wash publ (A): 1-279 pp.
- Costache, M.A.; Campeanu, G. and Neata, G. (2012). Studies concerning the extraction of chlorophyll and total Carotenoids from vegetables. *Romanian Biotechnology Letters*, 17(5): 7702-7708.
- Cowardiana, L.M.; Carter, V.; Golet, F.C.; La Roe, E.T. (1979). Classification of wetlands and Deepwater Habitats of the united states. FWS/OBS-79/31, U.S. Fish and wildlife service: Washington, D.C
- Crowder, A.A.; Bristow, J. M.; King, M.R. and Vander, K.S. (1977). Distribution, seasonality and biomass of aquatic macrophytes in lake Opnicon. *Naturaliste., Can.*, (104):441-456.
- Das, K. K. (2013). Diversity of aquatic and wetland angiospermic macrophytes in the Kamrup District, Assam, India. *Pleione*, 7(1): 8-17.

- Das, K.K. and Chakraborty, A.K. (2017). Elements of Biology, part-I, Published by Bina Library, Guwahati, Assam, India, 136-137 pp.
- Dar, M.N.; Manzoor, M. F.; Kaushik, V.; Kumar, M.; Rawat, S.; Shah, K.M.; Zaman, B. and Singh, E.A. (2017). "Water Quality Assessments of Dal Lake, Jammu & Kashmir". *International Journal of Scientific & Engineering Research*, 8(12): 328-337.
- Daubenmire, R.F. (1947). Plants and Environment A Textbook of Plant Autecology. Published by John Wiley & Sons, New York, 424 pp.
- Day, F. (1878). The Fishes of India: Being a natural history of the fishes Known to inhabit the seas and freshwater of India, Burma and Ceylon. Barnard Quaritch London. 778 pp.
- Day, F. S. (1978). The Fishes of India. William and Sons Ltd., London
- Day, F. (1885). Relationship of the Indian and African Freshwater fish fauna. *Journal of the Linnean Society of London, Zoology*, 18(107): 308-317.
- Day, F. (1889). The Fauna of British India, including Ceylon and Burma. I Taylor and Francis, London. 548 pp.
- De Silva, S.S. (1985). Performance of *Oreochromis niloticus* (L) fry maintained on mixed feeding schedules of different protein content. *Aquaculture and Fisheries Management*, 16: 331-34.
- De Silva, S.S.; Abery, N.W. and Nguyen, T.T.T. (2007). Endemic freshwater finfish of Asia: distribution and conservation status. *Diversity and Distributions*, 13: 172-184.
- De, A.K. (2016). Environmental Chemistry, 8<sup>th</sup> edition, published by New Age International Pvt. Ltd. 7/30 A, Darya Gang, New Delhi, India.
- Deb, S. and Singh, T.H. (2012). Status of Udali Beel in Karbi Anglong District, Assam. ISBN 978-81-202-8869-7. 70-79 pp.

- Debnath, D. and Yengkokpan, S. (2017). Fish feeds and self-sufficiency in fish production. The Assam Tribune (English Daily), June. 26, P.6, col. 2
- Deepa, P.; Raveen, R.; Venkatesan, P.; Arivoli, S. and Samuel, T. (2016). Seasonal variations of physicochemical parameters of korattur lake, Chennai, Tamil Nadu, India. *International Journal of Chemical Studies*, 4(3): 116-123.
- Deka, K. and Dutta, A. (2013). Ichthyofaunal Diversity and Status in Barbila Beel, Nalbari, Assam. *The Clarion: Multidisciplinary International Journal*, 2(2): 32-37.
- Deka, U. and Sarma, S.K. (2014). Present status of aquatic macrophytes of the wetlands of Nalbari district of Assam, India. *Asian Journal of Plant Science and Research*, 4(3): 67-75.
- Dembitsky,V. M.; Rozentsvet, O.A. and Zhuicova,U.S.(1992).Fatty acid and phospholipids from lichens of the order Leconorales. *Phytochemistry*. 31:.851-853.
- Denny,P.(1985). Wetland vegetation and associated plant life forms. In: the Ecology and Management of African wetland vegetation. P. Denny (ed.)- Dr Junk Publishers.1-18 pp.
- Deshmukh, S.B.; Phadke, N.S. and Kothandaraman,V.(1964). Physico-chemical characteristics of Ambazari lake Water. *Environmental Health*, 6: 186- 188.
- Devi, C.G. and Sharma, B.M. (2007). Studies on the diversity of the macrophytes in Awangsoipal Lake (Bishnupur), Manipur, India. In (Eds. A.K. Kandy and Asha Gupta) Biodiversity conservation and Legal aspect. Aviskar Publishers, Distributors, Jaipur. 62-71 pp.
- Devi, R.K. (2005). Investigation on physic- chemical condition of major effluents Points of Nambul river, Manipur, India. *Indian Journal of Environment and Ecoplannig*, 10 (2): 537-540.

- Devi, N.I.; Singh, N.R.; Devi, C.B. and Shomorendra, M.(2014). Ichthyofaunal Diversity of Utra Lake in Manipur. *International Journal of Current Research*, 6(12): 10815-10820.
- Devlin, R.M.and Witham, F. H. (1986). Plant physiology, Fourth Edition,CBS Publishers & Distributors, 485, Jain Bhawan,Bhola Nath Nagar,Delhi (India), 223-225 pp.
- Devlin, R.M. and Barker A.V.(1971).*Photosynthesis*, NewYork:VanNostrand, Reinhold.
- Dewanji, A. (1993). Amino acid composition of leaf proteins extracted from some aquatic weeds. *Journal of Agricultural and Food chemistry*, 41: 1232-1236.
- Dey, S.C. (1982). A critical Analysis on the fish and fisheries of Assam. *Proceding of National Seminar on Ichthyology*. 3:16.
- Dey, S.C. (1973). Studies on the Distribution and Taxonomy of the Ichthyofauna of the hill streams of Kamrup-Khasi-Garo Regions of Assam with special reference to the Functional morphology of some rheophilic Fishes, D.Sc. Thesis, University of Calcutta (India), xi +299 pp.
- Dey, S.C.(1981). Studies on the Hydrobiological Conditions of some Commercially Important Lakes (Beels) of Kamrup District of Assam and their Bearing on Fish Production. Final Technical Report, Northeastern Council, Govt. of India, Shilling, 1-177 pp.
- Dey, S.C.and Kar, D.(1989a).Aquatic macrophytes of Lake Sone in Assam. *Environment and Ecology*, 7(1), 253-254.
- Dey, S.C. and Kar, D. (1990). Fish yield trend in Sone, a tectonic lake of Assam. *Matsya*, 15-16: 39-43.
- Dey, S.C. and Kar, D. (1989). Aquatic Macrophytes of Lake Sone in Assam .*Environment and Ecology*, 7(1): 253-254.
- Dey, S.C. and Kar, D. (1989). Fishermen of Lake Sone in Assam; Their socio-economic status. *Science and Culture*, 55: 395-398.

Dhore, M.M. and Luchare, P.S. (2014). Survey of aquatic macrophytes in Yevatmal district, Maharashtra, India. *International Journal of Life Sciences*, 2(3): 273-275.

Dosunmu, M.I. (1997). Chemical composition of the fruit of *Tetrapleura tereptra* and the physico-chemical Properties of its oil. *Global Journal of Pure and Applied Sciences*, 3(1): 61-67.

Dutta, U. (2005).Ecological status of wetlands of Rudrasagar area in Sivasagar district, Assam: Ph.D. thesis. Gauhati University, Assam.

Dutta, T.; Deka, U. and Rabha, P.K. (2014). Diversity of aquatic macrophytes of Kapla beel (wetland) of Barpeta district, Assam, India. *Annals of Biological Research*, 5(12): 41-45.

Dutta, R.; Baruah, B.; Sarma, S.K.(2011). Influence of riparian flora on the river bank health of a Himalayan River before being regulated by a large dam in North East India. *Annals of Biological Research*, 2(4): 268-280.

Dutta, S.K. (2012). Wetlands of Assam: An overview. Published by Dr. S.K. Dutta, Principal / President, National Seminar Organizing Committee, Dhing College, Nagaon, Assam. 1-1 pp.

Edmondson,W.T.(1966).Changes in the oxygen deficit of Lake Washington, Internationale vereigung fur Theoretische und Angewandte Limnologie: Verhandlungen. 16(1):153-158.

Edwards, P.; Kamal, M.; Wee, K.L. (1985). Incorporation of composted and dried water *hyacinth* in pelleted feed for the tilapia, *Oreochromis niloticus* (Peters). *Aquaculture and Fisheries Management*. 16: 233-248.

Edwards, P. (1980).Food potential of Aquatic macrophytes.1<sup>st</sup> ed. International Center for living Aquatic Resources Management.

Edwards, P. (1987). Use of terrestrial and aquatic macrophytes in aquaculture. In: D. Moriarty and R.S.V. Pullin, Editors, Detritus and Microbial Ecology in Aquaculture. ICLARM conf. Proc. Vol. 14: 311-385 pp.

- FAO/WHO. (1984). Contaminates. In: Codex Aliment – arius (1<sup>st</sup> ed, XVII), FAO/ WHO. Codex Aliment –arius (Is ted,xvii) .FAO/WHO.Codex Alimentarius commission, Rome.
- Falge, R.; Schpanof, L. and Jurss, K (1978). Amylase, esterase and protease activity in the intestine content of rainbos trout salmo gaindneri rich after feeding with feed containing different amounts of starch and protein. *Journal of Ichthyology*, 48: 283- 287.
- Fisher, R.A. and Yates, F. (1963). Statistical Tables for Biological, Agricultural and Medical Research. 6<sup>th</sup> ed., Oliver & Boyd, Edinburgh and London, X, 146 pp.
- Fischer, Z. (1972). The elements of energy balance in grass carp (*Ctenopharyngodon idella* ) Part -III. Assimilability of proteins, Carbohydrates and lipids by fishseed with plant and animal food. *Pols. Arch. Hydrobiologia*. 19: 83-95.
- Fleek, H. (1981). Introduction of Nutrition. 4<sup>th</sup>. Edition. NewYork: MacMillan Publishing Company.
- Food and Agriculture organization (FAO) (2003). Food energy methods of analysis and conversion factors. *Food and nutrition paper*. Food and agriculture organization of the United states. 77: 1- 93 pp.
- Frenette, J.J.; Vincent, W.F.; Legendre, L. and Nagata, T. (1996). ‘Size dependent changes in phytoplankton C and N uptake in the dynamic mixed layer of Lake Biwa’. *Fresh Water Biology*, 36(2): 221-236.
- Fynn- Aikin, K.; Silas, S.O.; Hung, S.S.O. and Huges, S.G. (1993). Effects of feeding a fish high level D-glucose on liver function in Juvenile white sturgeon (*Acipenser transmontanus*). *Fish Physiology and Biochemistry*, 12: 317- 325.
- Game, A.S.and Salaskar, P.B.(2007). Environmental impact of macrophytes on Makhmali Lakes, Thane, Maharashtra. *Journal of Aquatic Biology*, 22(2): 203-204.

- Ganapati, S.V. (1973). Ecological Problems of man- made lakes of South India. *Archiv fur Hydrobiologie*, 71: 363-380.
- Gatlin III, DM and Wilson, RP. (1986). Dietary copper requirement of fingerling channel catfish. *Aquaculture*, 54: 277-285.
- Gayevskaya, N.S. (1966). The Role of Fresh water plants in Animal Nutrition *Nymphaeales*, Nanka, Moscow.1-327 pp.
- Gerloff, G.C.; Krombholz, P.H. (1966). Tissue analysis as a measure of nutrient availability for the growth of angiosperm aquatic plant. *Limnology and Oceanography*, 11: 529-539.
- Ghavzan, N.L.; Gunale, V.R.; Mahajan, D.M. and Shrike, D.R.(2006). Effect of environmental factors on ecology and distribution of aquatic macrophytes. *Asian journal of plant Sciences*, 5(5): 871-880.
- Ghosh, S.K. and Lipton, A.P. (1982). Ichthyofauna of the N.E.H. Region with special reference to their economic importance. ICAR Spl. Bulletin No. 1 (ICAR Research Complex, Shilling),119-126 pp.
- Ghosh, S.K. ( 2005). Illustrated aquatic and wetland plants in Harmony with Mankind, standard literature, Kolkata, 1- 225 pp.
- Ghosh, S.K. and Lipton, A.P. (1982). Ichthyofauna of the N.E.H. Region with special reference to their economic importance. ICAR Spl. Bulletin No. 1 (ICAR Research Complex, Shilling), 119-126 pp.
- Glass, B. (1961). Summary. In W.Mc Elroy and B.Glass, eds., *Light and Life*. Baltimore, Md : Johns Hopkins university press.
- Gogoi, R. (2006). Macrophytic Diversity of Wetlands of Deeper Beel with special references to seasonal variation. Ph.D. Thesis, Gauhati University, Guwahati, Assam
- Goodwin, T.W. (1958).Studies in carotenogenesis. 24. The changes in carotenoid and chlorophyll pigments in the leaves of deciduous trees during autumn necrosis. *Biochemical Journal*, 68(3): 503-511.

- Goodwin, T. (1960). Chemistry, biogenesis and physiology of the carotenoids. In W. Ruhland, ed., *Encyclopedia of plant physiology*. 5; part 1, 394, Berlin: Springer
- Goodwin, T.W. and Mercer, E.I (1983) 'Introduction to Plant Biochemistry'. 2<sup>nd</sup> ed., Pergamon Press, Oxford, 96-103 pp.
- Gopal, B. and Sah, M. (1995). Inventory and classification of wetlands in India. *Vegetatio*, 118: 39-48.
- Goswami, T. K. and Goswami,M.M.(2009). Freashwater Fishes of Jamlai wetland, in Kamrup District of Assam and its economic importance Fish and Fisheries in North East India. Recent Advances and Rebuilding Edited by Bhuyan,R,N.; Gosh,D. & Sarma,D. Ist Edition, 218-232 pp.
- Goswami, C.and Kalita, M.P. (2012). Ichthayofaunal Diversity and Anthropogenic stress on Deepor Beel: the only Ramsar site in Assam, IOSR *Journal of Environmental Science, Toxicology and Food Technology*, 2(1): 54-59.
- Gothberg, A.; Greger, M.and Bengtsson, B. (2002). Accumulation of heavy metals in water spinach (*Ipomoea aquatic*) cultivated in the Bangkok region, Thailand. *Environmental Toxicology and Chemistry*, 21: 1934-1939.
- Gowda, N.K.S.; Rammana, J.V.; Prasad, C.S. and Sing, K. (2004). Micronutrient content of certain tropical conventional and unconventional feed resources of southern India.*Tropical Animal Health and Production*, 36: 77-94.
- Gulmira S., Knapp M.andLichtenthaler H.K. (2006).Differences in photosynthetic activity, chlorophyll and carotenoid levels and in chlorophyll fluorescence parameters in green sun and shade leaves of *Ginkgo* and *Fagus*. *Journal of plant physiology*, 7:945-955

- Gupta, P.K. (2009). Methods in environmental Analysis water, soil and air. Published by AGROBIOS (India), Agro House, Behind Nasrani Cinema Chopasani Road, Jodhpur, India.
- Gupta, S. P. (2002). Statistical Methods. Published by Sultan. Chand & Sons, 23 Daryaganj, New Delhi.
- Hamilton, F.B. (1822). An account of the fishes found in the River Ganges and its branches. Archibald constable and Company, Edinbarg, London, 405pp, 39 pls.
- Hamilton, F. (1808-1814). *Vertebrates of Assam*, A Checklist with IUCN Status. Published by Assam Science Technology and Environment Council, Guwahati, Assam, India, 11 p.
- Hannah, M.A.C. and Krishnakumari, S.(2015). Analysis of Mineral elements, Proximate and Nutritive value in *Citrullus vulgaris* Schrad (Watermelon) seed extracts. *The pharma innovation Journal*, 4(8): 07-11.
- Hari, B.and Kurup, M.(2003). Comparative evaluation of dietary Protein levels and plant animal ratios in *Macrobrachium rosenbergii* (De. Mann.) *Aquaculture nutrition*, 9: 131- 137.
- Harney, N.V.; Dhamani, A. A. and Andrew,R.J.(2013).Biodiversity of Macrophytes of Three Water Bodies Near Bhadrawati, District-Chandrapur (M.S.), India. *International Journal of Scientific Research (IJSR)*, 2(9): 437-439.
- Harney, N.V. (2014). Macrophytes Biodiversity of Dudhala Lake of Bhadrawati, District-Chandrapur (Maharashtra State), India. *Asian Journal of Multidisciplinary Studies*, 2(4): 69-72.
- Hasan, M. R. and Chakrabati, R. (2009). Use of algae and aquatic macrophytes as feed in small scale aquaculture: a review; FAO, Fisheries and aquaculture, Technical paper No. 531. Rome, FAO, 123 p.
- Hastings, W.H. (1976). Fish nutrition and fish feed manufacture FAO Technical Conf. on *Aquaculture*, Kyoto, Japan, FIR: AQ/cont/76/R.23:13

- Haskin, H.H. (1942). A Spectrophotometric method for the analysis of Chloroplast Pigments. *Journal of Biological Chemistry*, 144: 149-160.
- Hertz, Y.; Madar, Z.; Hepper, B. and Gertler, A.(1989). Glucose metabolism in the common carp (*Cyprinus carpio* L: The effects of cobalt and Chromium. *Aquaculture* , 76: 255- 267.
- Hilton, J.W. (1989). The interaction of vitamins, minerals and diet composition in the diet of fish. *Aquaculture*, 79: 223-244.
- Hora, S. L. (1921a). Fish and fisheries of Manipur with some observations on those of Naga Hills. *Records of Indian Museum*, 22: 166-214.
- Hora, S.L (1941).The game fishes of India, XVI: The mahseers or the large-scaled barbs of India, 6: The‘Jungha’ of the Assamese *Barbus (Tor) progenius* Mc Clelland. *Journal of the Bombay Natural History Society*, 42: 526-532.
- Hosetti, B.B. (1995).“Treatment of Sugar industry effluents by ponds and lagoons”. *Journal of Environmental Biology*, 16: 143-149.
- Hosetti, B.B. and Kumar, A. (2001). A Text book of Applied Aquatic Biology .Published by Daya Publishing House, New Delhi
- Huheey, J.E.; Keiter E. A. and Keiter, R.L. (2001). Inorganic Chemistry: Principles of Structure and Reactivity, 3<sup>rd</sup> ed., Harper Collins College Publishers, New York.
- Hussainy, S.U. (1967). ‘Studies on limnology and Primary Production of a tropical lake; *Hydrologic*, 30: 335-352 pp.
- Hutchison, G.E. (1975). A treatise on Limnology, Chemical Compositions of *Ipomea aquatic* (Green Kangkong). *International Journal of Pharma and Bio Sciences*, 2: 594-598.
- Imhoff; Jahounes; Hans, F.; Sahl, G.; Gaber, S.H. Saliman, and Hans, Trueper, G. (1980).“The wadi naturn, Egypt. Chemical composition and microbial mass developments in alkaline brines of eutrophiobic desert lakes”. *Geomicrobial Journal*, 1(3): 219-234.

- Ipsen, J. and Feigl, P. (1970). *Bancroft's introduction to biostatistics*. 2<sup>nd</sup> ed., Harper and Row Publishers, New York, Evanston and London
- Islam, Z. and Rahmani, A.R. (2008). Potential and Existing Ramsar Sites in India. Oxford University Press, Oxford, New York, 1-578 pp.
- Islam, M.S.; Ismail, B.S.; Barzani, G. M.; Sahibin, A. R. and Ekhwan, T.M. (2012). "Hydrological Assessment and Water Quality Characteristics of Chine Lake, Pahang, Malaysia". *American-Eurasian Journal Agricultural and Environmental Sciences*, 12 (6): 737-749.
- IUCN (2011). IUCN Red list of Threatened species. Version [www.iucnredlist.org](http://www.iucnredlist.org), Downloaded on, 19 November, 2016.
- Jain, V.K. (2007). Fundamentals of plant physiology. S. Chand & company Ltd ., Ram Nagar, New Delhi. 186 p.
- James, C.S. (1995). Analytical Chemistry of food, 1<sup>st</sup> edition, Chapman & Hall. 1, 56- 57 pp.
- Jayaram, M.C. and Shetty, H.P.C. (1980). Studies on the growth rates of catla, rohu and common carp fed on different formulated feeds. *Mysore Journal of Agricultural Sciences*, 14(4): 598-606.
- Jayaram, K.C. (1981). The freshwater fishes of India, Pakistan, Bangladesh, Burma and Sri Lanka-A Handbook Zoological Survey of India, Calcutta, 1- 475 pp.
- Jayaram, K.C. (1999). The Freshwater Fishes of the Indian Region. Narendra Publishing House, Ned Delhi, 1-322 pp.
- Jayaram, K.C. (2010). Fishesh of the Indian Region, Narendra Publishing House (New Delhi).
- Jeffrey, S.W. and Mantoura, R.F.C. and Wright, S. W. (1997). Phytoplankton pigments in oceanography: guidelines to modern methods. UNESCO Monographs on Oceanographic Methodology, UNESCO Publishing, Paris.10

- Jhingram, V. G. (1992). Fish and Fisheries of India, 3<sup>rd</sup> Edn., Hindustan Publishing Corporation ( India) Delhi. 463pp.
- Jiang, S. L.; Wu, J.G.; Feng,Y.;Yang,X.E. and Shi,C.H.(2007). Correlation Analysis of Mineral Element Contents and Quality Traits in Miled Rice (*Oryza sativa l.*). *Journal of Agricultural and Food Chemistry*, 55 (23): 9608-9613.
- Kachroo, P. (1984). Aquatic Biology in India. Bishen Singh Mahendra Pal Singh, Dehra Dun, India.
- Kalita, B.; Bhuyan, K.C.and Dutta, A. (2000). Studies on Fishery Potential of the river Sonai of Morigaon District, Assam. *Journal of Assam Science Society*, 41 (3): 173 -185.
- Kalita, B.; Bhuyan, K.C.; Kusre, D. and Dutta, A. (2006). Physico- chemical quality of Beel water in Morigaon district, Assam. *Journal Ecobiology*, 18(10): 17-21.
- Kalita, P.; Mukhopadhyay, P.K.; Mukherjee, A. K. (2007). Evaluation of the nutritional quality of four unexplored aquatic weeds from northeast India for the formulation of cost-effective fish feed. *Food Chemistry*, 103: 204-209.
- Kalita, G. (2008). Ecology and Distribution of macroinvertebrate enmeshed fauna in Deepar wetland of Assam, India. Ph.D. Thesis of Gauhati University, Guwahati, Assam, India.
- Kalita, S.; Kalita, H.; Das, M.; Saikia, M.D. (2018). Assesssment of Spatial Variations of Water Quality Index of Deepor Beel, Assam, India. *International Research Journal of Engineering and Technology (IRJET)*. 05 (06): 775-780.
- Kalita, P.; Pathak, S.; Deka, P. (2016). A Preliminary Study on Ichthyofaunal Resource of Motapung-Maguri Beel of Tinsukia District of Assam, India. *International Journal of Fauna and Biological Studies*, 3(4): 97-102.

- Kalita, S. R. and Choudhury, K. (2016). Aquatic Macrophytic Diversity of Upod Beel, Goalpara District of Assam, India. *Asian Resonance*, 5(4): 87-91.
- Kalita, G. and Sarma, S.K. (2019). Aquatic Macrophytic and Ecotone floral diversity, species distribution analysis in certain wetlands (Beels) of Kamrup district of Assam. *International Journal of Basic and Applied Research*, 9(6): 670-684.
- Kalla, A.; Bhatnagar, A.; Garg, S.K. (2004). Further studies on protein requirements of growing Indian major carps under field conditions. *Asian Fisheries Science*, 17: 191-200.
- Kannan, K. (1991). Fundamentals of Environmental Pollutions, S. Chand and Company Ltd., New Delhi, 189 pp.
- Kapoor, D.; Dayal, R. and Ponniah, A.G. (2002). Fish biodiversity of India. NBFGR, Lucknow, India Publication, 775 pp.
- Kar, D. and Barbhuiya, M. (2000). Studies on the aquatic macrophytes of Chatla Haor, flood- plain wetland in Cachar district in Assam. *Indian Science Congress*, New Delhi.
- Kar, D. (1984). Limnology and Fisheries of Lake Sone in Cachar district of Assam, India. Unpublished Ph.D.Thesis, Gauhati University, Assam, 201 pp.
- Kar, D.(2003). Fishes of Barak drainage, Mizoram and Tripura, Pp. 203-211. In: Kumar, A.C. Bora and L.K. Singh (Eds.). Environment Pollution and Management, APH Publishing Corporation, New Delhi, 1-604 pp.
- Kar, D. and Dey, S.C.(1993). Inter relationship and dynamics of fish population of lakes Sone in Assam. *Environment Ecology*, 11(3):718-719.
- Kar, D. and Dey S.C. (2000). Yield and Conservation of Indian major Carps of Lake Sone in Assam. *Environment and Ecology*, 18(4), 1036-1038.

- Kar, D. and Barbhuiya, M.H. (2000b). Ichthyodiversity of Chatla Haor: a floodplain wetland in Barak valley region of Assam, 3-6, in *Advances in Zoology and Environmental Degradation and Biodiversity*, (eds) Pandey B.N. and Singh B.K., Daya Publishing House, New Delhi, 279 pp.
- Kar, D.; Kumar, A.; Bohra, C. and Singh, L.K. (2003). Fishes of Barak drainage, Mizoram and Tripura. In: Environment, Pollution and Management, APH Publishing Corporation, New Delhi, 604: 203-211.
- Kar, D. (2006). Fish Fauna of River Barak of Mizoram and of Tripura with a note on Conservation. *Journal of Freshwater Biology*, 16.
- Kashiwada, K.; Teshima, S. and Kanazawa, A. (1970). Studies on the production of B vitamins by intestinal bacteria of fish-V. Evidence of the production of Vitamin B<sub>12</sub> by microorganism in the intestinal canal of carp *Cyprinus Carpio*. *Nippon Suisan Gakkaishi*, 36: 421- 424.
- Kashyap, D.; Hazarika, D.; Bhattacharjya, B.K.; Bhuyan, S.; Phukan, B.; Baishya, S. and Bordoloi, R. (2012). Ichthyofaunistic Diversity of Selected Floodplain Wetlands of Central Brahmaputra Valley (CBV) Zone of Assam With a Note on Their Conservation. In. Current issues of conservation and wise use of wetlands in North Eastern Region of India. Published by Dr. S.K. Dutta, Principal / President National Seminar (UGC Sponsored National Seminar Proceedings, 3 & 4 February, 2012) Organizing Committee, Dhing College, Nagaon, Assam. 128-140 pp.
- Kaul, V.; Trishal, C.L. and Handoo, J.K. (1980). Distribution and production of macrophytes in some water bodies of Kashmir In: Singh J.S., and Gopal B. (eds.), *Glimpses of Ecology*, Prof. R. Mishra Commemoration volume, 313-334 pp.
- Kaushik, S. and Saksena, D.N. (1999). Physico- chemical limnology of certain water bodies of central India. Pages 1- 58. In. Vijaykumar, k. *Freshwater ecosystem of India*. Daya Publishing House, Delhi, 1- 336 pp.

- Keddy, P.A. (2000). Wetland Ecology: Principles and Conservation, Cambridge University press, Cambridge.
- Khalil, E.; Esoh, R.; Rababah, T.; Almajwal, A.M. and Alu'datt, M.H. (2012). Minerals, Proximate Composition and their correlations of medicinal plants from Jordan. *Journal of Medicinal Plants Research*, 6(47):5757-5762.
- Khan, M.A. (1987). Observation of zooplankton composition, abundance and periodicity in two floodplain lakes of Kashmir valley. *Acta hydrochem. Hydrobiol.*, 15: 166-174.
- Khan, M.A.; Jafri, A.K.; Anwar, M.F.; Hasan, M.A. and Erfanullahh, M. (1991). Nutritional efficiency of oilcakes in diets compounded for fingerlings of Indian major carp *Labeo rhabta* (Ham.). Proc. of National symposium on Horizons in Freshwater Aquaculture, 23-25 January, 73-77 pp.
- Khan, M.A.G. and Choudhary, S.H. (1994). Physical and Chemical limnology of Lake Kaptai, Bangladesh. *Tropical Ecology*, 35(1): 35-51.
- Khan, K.Y.; Khan; M.A.; Niamat, R.; Munir, M.; Fazal, H.; Mazari, P.; Seema, N.; Bashir, T.; Kanwal, A. and Ahmed, S.N. (2011). Element content analysis of plants of genus *Ficus* using atomic absorption spectrometer. *African Journal of Pharmacy and Pharmacology*, 5(3): 317-321
- Khan, R.M.; Jadhav, M. J.; Ustad, I.R. (2012). "Physicochemical Analysis of Triveni Lake Water of Amravati District in (M.S.) India. *Bioscience Discovery*, 3(1): 64-66.
- Kiran, B.R.; Patel, A.N.; Vijaya, K. and Puttaiah, E.T. (2006). Aquatic macrophytes in fish culture ponds at Bhadra fish farm, Karnataka. *Journal of Aquatic Biology*, 21(2): 107-130.
- Kodarkar, M.S. (1996). Conservation of lakes, case study of five lakes around Hyderabad, Andhra Pradesh, India Indian Association of Aquatic Biologist (IAAB) 3<sup>rd</sup> publication. Hyderabad, 85 pp.

- Kono, M.; Matsuinal and Shimizol (1987). 'Effect of Chitin, Chitson and cellulose as diet supplementary in the growth of culture fish'. *Bull. Top. Soc. Fish.* 53(1): 125- 129.
- Kosygin, L.; Dhamendra, H.and Gyaneshwari, R. (2007). Pollution status and conservation strategies of Moirang river, Manipur with a note on its aquatic bio-resources. *Journal of Environmental Biology*, 28(3): 669- 673.
- Kosygin, L. and Dhamendra, H. ( 2009). Ecology and Conservation of Loktak Lake, Manipur. An overview. In: Wetlands of North East India. Ecology, Aquatic Bioresources and conservation (Eds; L. Kosygin). Akansha Publishing House, New Delhi, 1-20 pp.
- Kottelat, M. and Whitten, T. (1996). Freshwater biodiversity in Asia with special reference to fish: World Bank Technical Paper, 343:59.The world Bank, Washington, D.C., 17-22 pp.
- Kumar, S. and Sharma L. (1991). "Comparative physico-chemical limnology of lakes Pichhola and Fatehsagar, Udaipur (Rajasthan)".*Pollution Research*, 10 (3): 173-178.
- Kumar, S.(1997). Limnological studies on Kunjwani Pond, Jammu. 1. Physicochemical factors and its scope to fish culture. Recent advances in fresh water biology, Vol. -11. Anmol Publication, New Delhi, 1-20 pp.
- Kumar, R.and Pandit, A.K. (2005). Community architecture of macrophytes in Hokarsar wetland, Kashmir, India. *Indian Journal of Environment and Ecoplanning*, (10): 565-573.
- Kumar, A. (2014). Studies on Ichthyofaunal Diversity with special reference to monthly and seasonal variations of fish landings in glacial fed mountainous Goriganga River of Kumaun Himalaya, Uttarakhand, India. *Research Journal of Animal, Veterinary and Fishery Sciences*, 2(4): 1-2.

- Lahon, B. (1983). Limnology and fisheries of some commercial beels of Assam, India. Unpublished Ph. D. Thesis, Gauhati University, Assam, 349 pp.
- Laishram, J. and Dey, M. (2014). Water quality status of Loktak lake, Manipur, North east India and Need for conservation Measures: A study of five selected villages. *International Journal of Scientific and Research Publication*, 4 (6): 1-5.
- Lakshmi, B.B.; Naidu,Y. P. and Rao,K.R.(2015). Ichthyo Faunal Diversity and Conservation of Kolleru Lake- A Ramsar site in Andhra Pradesh. *IOSR Journal of Pharmacy and Biological Sciences*. 10(2): 13-21.
- Lall, S.P. (1989).The minerals. In: J.E. Halver (Editor), Fish Nutrition, 2<sup>nd</sup> edn. Academic Press, New York, 219-257 pp.
- Larson, G.L.; McIntire, C.D.; Karnaugh Thomas, E. and Hawkins Haffiman, C. (1995). "Limnology of isolated and connected high-mountain lakes in Olympic National Park, Washington State, U.S.A." *Archiv fuer Hydrobiologia*, 134(1): 75-92.
- Lavania, G.S.; Paliwal, S.C and Gopal, B. (1990). Aquatic vegetation of the Indian subcontinent. In :(Ed.) Brij Gopal, Ecology and Management of Aquatic vegetation in the Indian subcontinent. Kluwer Academic Publishers, Dordsecht Boston/ London, 29-76 pp.
- Latif, K.A.; Alam, M.T.; Sayeed, M.A.;Hussain, M.A.;Sultana, S.and Hossain, M.A.(2008). Comparative study on the effect of low cost oil seed cake and fish meal as dietary protein sources for *Labeo rohita* (Hamilton) fingerlings. *University Journal of Zoology, Rajshahi. Univiversity*, 27: 25-30.
- Lehr, J.H.; Tyler, E.G.; Wayne, A.P. and Jack, D. (1980). Domestic water Treatment. New York; Mc Graw Hill Book Company.
- Leveque, C.; Oberdorff, T.; Paugy, D.; Stiassny, M.L.J. and Tedesco, P.A. (2008). Global diversity of fish (Pisces) in freshwater. *Hydrobiologia*, 595: 545-567.

- Lichtenthaler, H. K. (1987)."Chlorophylls and Carotenoids: Pigments of Photosynthetic membrane", Methods in Enzymology, Academic Press, New York, Vol. 148: 350-382 pp.
- Lim, R.B.L.; Tan, H.T.W. and Chua, K.S. (1998). Freshwater macrophytes. In: P.K.L. Ng(Ed.), A Guide to Freshwater life in Singapore, BP Singapore Pvt. Ltd. and Omni- Theatre, Singapore Science Centre.
- Limsuwan, T. and Lovell, R.T. (1981). Intestinal synthesis and absorption of Vitamin B<sub>12</sub> in Channel Catfish. *Journal of Nutrition*, 111: 2125-2132.
- Lovell, R.T. (1972).'Protein requirement of cage cultured channel catfish'. Proc.26<sup>th</sup> Ann. Conf. S. *East Asian Game Fish. Comm*, 357-361 pp.
- Lovell, R.T. (1979). Formulating diets for aquaculture species. *Feedstuffs*, 51: 29-32.
- Lovell, R.T. (1989).Nutrition and Feeding of Fish van Mostrand Reinhold, New York, 260 pp.
- Maheshwari, J.K.(1960). The Origin and Distribution of the naturalized plants of Khandwa Plateau, Madhya Pradesh. *Journal of Botanical Science*, 3: 9-19.
- Maibam, S.; Ngasepam, R.S.and Ningthoukhongiam, I.D. (2015). Check list of Fish Species of Loktak Lake, Bishnupur District. *Biological Forum- An International Journal*, 7(1): 171-179.
- Majumdar, N.C. (1965). Aquatic and Semiaquatic flora of Calcutta and adjacent localities. *Bulletin Botanicai Society of Bengal*, 19: 10-17.
- Malakar, N.C. (1995). The systematic studies on the aquatic angiosperm of Cachar district of Assam, Ph. D. Thesis, Gauhati University .
- Manjare, S.A.; Vhanalakar, S.A.and Muley, D.V. (2010). Analysis of water quality using Physico-chemical Parameters Tamdalge tank in Kolhapur district, Maharashtra. *International Journal of Advanced Biotechnology and Research*, 1(2): 115-11 9.

- Mason, C.F. (1989). *Biology of Fresh of Fresh water Pollution*. U.K., Essex.: Longman Scientific and Tech. Publ.
- Meetei, W.S. and Singh, N.I. (2011). Effects of solid waste disposal on water in Imphal City, Manipur. *Pollution Research*, 30(1): 21-25.
- Menon, A.G.K. (1999). Checklist-freshwater fishes of India. Records of the Zoological Survey of India, Miscellaneous Publications, Occasional Paper No. 175:1-366.
- Meshram, C.B. and Dhande, R.R. (2000). Algae diversity with respect to pollution status of Wadali Lake, Amravati, Maharashtra, India. *Journal of Aquatic Biology*, 15(1&2): 1-5.
- Meshram C.B. (2003). Macro-invertebrate fauna of Lake Wadali, Amravati, Maharashtra. *Journal of Aquatic Biology*, 18(2):47-50.
- Michael, P.C. and Nicholas, A.W. (1998). Pigment composition of putatively achlorophyllous angiosperms. *Plant systematic, Evolution* 210: 105-111.
- Millennium Ecosystem Assessment (MEA) (2005). Ecosystems and Human Well-being Wetland: Wetlands and Water Synthesis. World Resources Institute, Washington, DC.
- Mir, A.M. and Kachroo, P.(1981).‘Limnology of Kashmir Lakes.VIII. Ecology of Bacillariophyceae in two lakes of Srinagar’. Proc. Indian Natn. Sci. Acad. Pt. B. 48(30): 378-390.
- Mirashi, M.V. (1954). Studies in the hydrophytes of Nagpur I. A preliminary survey. *Journal of Indian Botanical Society*, 33: 299-308.
- Mishra, S. R. and Sanksena, D.N. (2009). *Aquatic Ecology* Published by S.B. Nangia. A.P.H. Publishing Corporation. 4435-36/7,Ansari Road, Darya Ganj, New Delhi, 177-187 pp. and 275- 292 pp.
- Mitsch, W.J.; Gosselink, J.G.(1993). *Wetlands*. 2<sup>nd</sup> Ed. John Wiley and Sons (formerly Van Nostrand Reinhold), New York, 722 pp.

- Mittermeier, R.A.and Mitemeir,C.G.(1997). Megadiversity Earth biologically Wealthiest Nation. In mc Allister, D.E.A. Lttamiltion and B.Harvey (Eds). Global fresh water Biodiversity sea wind cemex, mexico city, 1-140 pp.
- Mohanty, S.N.and Swamy, D.N. (1986). Enriched conventional feed for Indian major carps. Proceedings of the First Asian Fisheries Forum, Manila, Philipphines, 26-31pp.
- Mohanty, S.N.; Swamy, D.N. and Tripathi, S.D. (1990).Growth nutritional Indices and carcass composition of Indian major carp fry *Catla Catla*, *Labeo rohita* and *cirrhinus mrigala* fed four dietary protein levels'. *Aquaculture Hungarica*, 6: 211- 217 pp.
- Molot, L.A. and Dillon, P.J. (1991). "Nitrogen: Phosphorus ratio and the prediction of chlorophyll in phosphorus limited lakes in Central Ontario". *Canadian Journal Fisheries and Aquatic Science*, 48: 140-145.
- Moore, T.C. (1981) "Research Experiences in Plant Physiology- A Laboratory Manual", Second Edition, Springer-verlag, New York, 35-4
- Mondal, T. K. and Ray, A. K. (1999). The nutritive value of *Acacia auriculiformis* leaf meal in compounded diets for *Labeo rohita* fingerlings. The fourth Indian Fisheries Forum Proceedings 1996, Kochi, 295-298 pp.
- Moynard, A.L. and Loosli, K.J. (1969). Animal nutrition. 5<sup>th</sup> Edition, Mc Graw Hill, New York.
- Munawar M. (1970). 'Immunological studies of fresh water Pond of Hyderabad, India-I'. *The Biotope, Hydrobiologia*, 3(5): 127-162.
- Murkute, V.B., and Chavan, A.W. (2016). Macrophytes diversity of three freshwater pond at Brahmapur Chandrapur (MS) India, *International Journal of researches in Biosciences, Agriculture and Technology*, 108-111pp.

- Muta Harah, Z.; Japar sidik, B.; Raesah, A. Maini, C. and Suzalina, A. (2005). Aquatic macrophytes in natural and manmade water bodies. *Bio-Science Research Bulletin*, 21(1): 27-36.
- Muztar, A.J., Slinger, S. J.and Burton, J.H. (1979). Chemical composition of Aquatic macrophytes.IV. Carotenoids, soluble sugars and starch in relation to their pigmenting and ensiling potential. *Canadian Journal Plant Science*, 59: 1093-1098.
- Myers, N. (1988).1.Threatened Biotas:“Hotspots” in tropical forests, *Environmentalist*, 8(3): 1-20.
- N.A.S. (National Academy of Sciences) (1984). Making aquatic weeds useful: some perspectives for developing countries. Washington, DC.
- Nadoni, M.B.; Murthy, P.S. and Hosetti, B.B. (2000). Water quality index of Amani tank. Tumkur, Karnataka state London: IAWAQ (in press)
- Nag, R.; Singha, N. and Deka, P. (2017). A Study on the Fish Diversity of Dhir Beel of Dhubri District of Assam, India. *International Journal of Applied Research*, 3 (5): 19 -26.
- Narayana, J.; Purushothama, R.; Kiran, B.R.; Ravindrakumar, K.P. and Puttah ,E.T.(2005). Investigation of drinking water quality of Basavanahole Tank with reference to physical Chemical Characteristics. *Fundamental of Limnology*, 201-206 pp.
- Narzari, S. and Sarmah, J. (2015a). A study on the prevalence of entomophagy among the Bodos of Assam. *Journal of Entomology and Zoology studies*, 3: 315–320.
- Narzari, S. and Sharmah, J. (2015b). Proximate composition of wild edible insects consumed by the Bodo tribe of Assam, India. *International Journal of Bioassays*, 4: 4050-4054.
- Nath, B. and Deka, C. (2012).A study on fish diversity, conservation status and anthropogenic stress of Chandubi tectonic lake, Assam, India. *Journal of Bio Innovation*, 1(6):148-155.

- Nath, P. and Dey, S.C. (1997). Fish and Fisheries of North-East India, Vol.1. Arunachal Pradesh, 140 pp.
- Nath, P. and Dey, S.C. (1989). Fish and Fisheries of North East India. Vol. 1: 1-143.
- Nath, P. and Dey, S.C. (2000). Fish and fisheries of North Eastern India (Arunachal Pradesh). Narendra Publishing house, Delhi. 217 pp.
- NationalWetlandAtlas: Assam (2010).SAC/RESA/AFEG/NWIA/ATLAS/18/2010, Space Applications Centre (ISRO), Ahmedabad, India,174p.
- Nayak, T.R. (1993)."Phytoplankton and certain physic- chemical conditions in freshwater Lake Matyata, Panna (M.P.) India".*Proceeding Academy Environmental Biology*, 2(2): 179-185.
- Nayaka, B.M.S. (2018).Ichthyofaunal Diversity of Mallasandra Lake of Tumakuru, Karnataka State, India. *Natural Journal of Multidisciplinary Research and Development*, 3(2):15-17
- Nayek, S.; Choudhury, I.H.; Nishika, J.and Roy, S. (2014). Spectrophotometric Analysis of Chlorophylls and Carotenoids from Commonly Grown Fern Species by Using Various Extracting Solvents. *Research Journal of Chemical Sciences*, 4(9): 63-69.
- NBSAP (2005). National Biodiversity Strategy and Action Plan, Prepared by the NBSAP Technical and policy core group, Kalpvriksh, Delhi.
- Nelson, J.S. (2006). Fishes of the world, 4<sup>th</sup> edition, John Wiley and Sons, inc, 601 pp.
- NRC (National Research Council) (1983). Nutrient Requirement of Warmwater Fishes and Shellfishes. National Academy Press, Washington, DC, 1-102 pp.
- Odum, E.P. (1971). Fundamentals of Ecology. W.B. Saunders Company, Philadelphia. 624 pp.

- Ogino, C. and Chen, M.S. (1973). Protein nutrition in fish, III. Apparent and true digestibility of dietary protein in Carp. *Bulletin Tropical Society of Fish.*, 39: 649-657.
- Olila, O.G.; Reddy, K.R. and Harris, W.G. (1995)."Forms and distribution of inorganic phosphorus in the sediments of two shallow eutrophic lakes in Florida". *Hydrobiologia* 302: 147-161.
- Omojola, A.B.; Uzohou and Fagbuar, S.S. (2012). Nutritive value and Potential of Aquatic plant (*Nephrolepsis bisserata* Schott) as feed resource for small ruminant. *Journal of Animal Science Advances*, 2: 749-754.
- Palit, D.; Kar, D. and Mukherjee, A.(2017). Studies on Macrophyte Diversity in Lalbandh Wetland, Birbhum, West Bengal, India. *International Journal of Interdisciplinary and Multidisciplinary Studies*, 4(2): 198-203.
- Pancho, J.V. and Soerjani, M. (1978). Aquatic weeds of Southeast Asia : A Systematic account of common Southeast Asian aquatic weeds. National Publishing Cooperative Incorporated, Philippines.
- Pandey, V.C. and Pandey, D.C. (1980). "Diatom flora of Allahabad (India)-1". *Proceeding of Indian National Science Academy*, B 46(3): 350-355.
- Pandey, B.N.; Das, P.K.L.; Dubey, S.V. and Hussain, S. (1999). Biomonitoring of water quality of river Ramjan (at Kishanganj) in relation to its impact on biological components pages: 310 – 336 in Vijay Kumar, K. Fresh water ecosystem of India. Daya Publishing House, Delhi, 1- 336 pp.
- Pandey, S.N and Sinha, B.K.(2013).Plant Physiology, Fourth Edition, Vikas Publishing House Pvt. Ltd., Noida, New Delhi, 1-229 pp.
- Pandit, A.K. (1984). Role of macrophytes in the aquatic ecosystems and management of freshwater resources. *Journal of Environmental Management*, 18(1): 73-78.

- Patra, S.S.; Mukhopadhy, A.K. and Saha, C. (2000). Effect of *Nymphoides* leaf meal incorporated diets on growth and body composition in *Cirrhinus mrigala*. In: the fifth Indian fisheries forum (Pp.77-79), India. Asian Fisheries Society, Indian branch, Mangalore.
- Pathak, N.B. and Mankodi, P.C. (2013). Hydrobiological status of Danteshwar pond, Vadodara, Gujarat, India. *International Research Journal Environmental Science*, 2(1):43-48.
- Pathak, H.; Pathak, D. and Limaye, S. N. (2012). Studies on the Physico-chemical status of two water bodies at Sagar city under anthropogenic influences. *Advances in Applied science Research*. 3(1): 31-44.
- Patnaik, S and Das, K.M. (1979). " Utilisation of some aquatic weeds as feed for rearing carp spawn and fry".*Sympo inland Aquaculture* 1977, Central inland Fishery Resources, Colombo (Sri Lanka). 40-416 pp.
- Pattar, M.; Teerthe, S.; Kerur, B.R. (2017). A study on major and trace elements in some traditional medicinal plants using AAS technique. *Int. Journal of Research in pharmacy and pharmaceutical Sciences*, 2(6): 86-90.
- Pattanaik, H. and Pattanaik, N.K. (1956). The Hydrophytes of Cuttack. *Journal of Indian Botanical Society*, 35: 167-170.
- Pawar, S.K.; Madlapur, V. R. and Pulle, J.S. (2003). Study on fish diversity in the Sirur Dam Near Mukhed Nanded (M.S.). *Journal of Aquatic Biology*, 18(2):60-70.
- Pejaver, M.; and Gurav, M. (2008). "Study of Water Quality of Jail and Kalwa Lake, Thane, Maharashtra".*Journal of Aquatic Biology*, 23(2): 44-50.
- Piecaynska, E. and Ozimek, T. (1976). Ecological Significance of Macrophytes. *International Journal of Ecology and Environmental Science*, 2:155-158.
- Pike, R.L. and Brown, M.L. (1967). Nutrition: an integrated approach. John Wiley and Sons. New York, 1- 542 pp.

Porra, R.J.; Thompson, W. A. and Kreidemann, P.E. (1989).Determination of accurate extinction coefficients and simultaneous equation for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectrometry, *Biochimica et Biophysica Acta*, 975: 384-394.

Porra, R.J. (1991).Recent advances and re-assessments in chlorophyll extraction and assay procedures for terrestrial, aquatic and marine organisms, including recalcitrant algae. In: Scheer H (ed.) Chlorophylls, 31-57.

Porra, R.J. (2002).The chequered history of the development and use of simultaneous equations for the accurate determination of chlorophyll a and b, *Photosynthesis Research*, 73: 149-156.

Pradhan, V.; Mohsin, M. and Gaikwad, B.H. (2012)."Assessment of Physico Chemical parameters of Chilika Lake water". *International journal of Research in Environmental Science and Technology*, 2(4): 101-103.

Prasad, S.N.; Ramachandra, T.V.; Ahalya, N.;Sengupta,T.; Kumar,A; Tiwari, A.K.; Vijayan, V.S.; Vijayan, L.(2002).Conservation of Wetlands of India- a review. *Tropical Ecology*, 43(1): 173-186.

Prasad, N. K.; Shivamurthy, G.R. and Aradhya, S.M. (2008). *Ipomoea aquatic*, An Underutilized Green Leafy Vegetables: A Review, *International Journal of Botany*, 4(1):123-129.

Prasad and Das (2018): Diversity and distribution of aquatic macrophytes with special reference to invasive species in Barak valley, Assam, Northeast India. *An international Journal of environment and biodiversity*, 9(1): 102-108

Puri, P. J.; Yenkie, M.K.N.; Sangal, S.P.; Gandhare, N. V.; Sarote, G.B. and Dhanorkar, D.B. (2011). "Surface water (Lakes) quality assessment in Nagpur city (India) based on Water quality index (WQI)". 4(1): 43-48.

- Purohit, R. and Singh, S.P. (1981). "Seasonal variation in physic-chemical limnology of shallow zones of Nainital Lake, Western Himalayas, India". *Proceeding of Indian National Science Academy B*, 47(2): 194-203.
- Puste, A.; Sarkar, P.K.; Das, D.K. (2004). Wetland ecosystem-a study on fish-aquatic food crops diversity for enhancing productivity and economic stability for fish-farm families in Indian subtropics. IIFET 2004 Japan proceedings.
- Puwastein, P.; Sion, T.E.; Craven, G.; Feliciano, R.R. and Judprasong, K.(2011). ASEAN Manual of Food Analysis, 1<sup>st</sup> ed. *Regional Centre of ASEAN networkof Food Data System*. Institute of Nutrition, Mahidol University, Thailand, 41-80 pp.
- Radhakrishnan, K.V. and Kurup, B.M.(2010). Ichthyodiversity of Periyar Tiger Reserve, Kerala, India. *Journal of Threatened Taxa*, 2(10): 1192-1198.
- Rahman,W.; Deka,R.; Kalita, B.and Deka, P.A. (2012). A Comparative Study on Ichthyofaunal Resource of Charan and Manaha Beel nof Morigaon District of Assam, India. *Int. J. of Fish and Aquatic Studies*, 4(4): 43-51.
- Rahman, A.H.; Rafiul, I.; Naderuzzaman, A.K.M.; Hossain, M.D. and Rowshatul, A. (2007). Studies on the Aquatic Angiosperms of the Rajshahi University Campus. *Research Journal of Agriculture and Biological Sciences*, 3(5): 474-480.
- Rai, D.N. and Dutta Munshi, J.M. (1988). Ecological characteristics of Chaurs of North Bihar. In; wetlands-Ecology and Management, vol. II (Gopal B., Turner R.E., Wetzel R.G. and Wingham D.F. eds.). International Science Publication and National Institute of Ecology, Jaipur, India. 88-95 pp.
- Rai, H. and Hill, G. (1982). On the nature of ecological cycle of Lago, January: A central Amazonian Rie-Verzea lake. *Tropical Ecology*, 23: 1-49.

- Ramakrishna Rao, S. (1990). 'Bioenergetics and methodology of energy flow in important wetlands and lake systems of Eastern Ghats.' *Env. Research*, New Delhi: Ministry of Environment & Forest Publication, 1-42 pp.
- Ramtake, D.S.; Moghi, C.A. and Mogh, C.A. (1989). Manual on water and waste water Analysis, National Environmental Engineering Research Institute (NEERI), Nehru Marg, Nagpur.
- Rashmi, C.L.; Sahana, S.M.; Sandesh, N. K.; Shivakumar, V.B. and Poornima, K.B. (2017). Water quality Analysis of Shantisagar Lake, Karnataka, India. *International Journal of Engineering Science and Computing*. 7(4): 10833-10838.
- Rath, R.K. (2000). Freshwater Aquaculture. 2<sup>nd</sup> edn. Jodhpur. Scientific Publications India.
- Ray, A.K.; Das, I. (1996). Evaluation of Dried Aquatic Weed, *Pistia stratiotes*, Meal as a Feedstuff in Pelleted Feed for Rohu, *Labeo rohita*, Fingerlings. *Journal of Applied Aquaculture*, 5(4): 35-44.
- Renukaradhya, K.M. and Verghese, T.J. (1986). 'Protein requirement of the carps, *Catla, catla* (Ham) and *Lebeo rohita* (Ham), Proc Indian Academy of sciences (Anim. Sci.) 95: 103- 107 pp.
- Ritchie, R.J. (2006). Consistent Sets of Spectrophotometric Chlorophyll Equations for Acetone, Methanol and Ethanol Solvents. *Photosynthesis Research*, 89(1): 27-41.
- Roginski, E.E. and Mertz, W. (1977). A Biphasic Response of Rats to Cobalt. *The Journal of Nutrition*, 107(8): 1537-1542.
- Romo, S. and Miracle, M.R. (1994)."Population dynamics and ecology of subdominant phytoplankton species in a shallow hypertrophic lake (Albufera of Valencia, Spain)". *Hydrobiologia*, 273(1): 37-56.
- Rowan, K.S. (1989). Photosynthetic pigments of algae. Cambridge University Press, Cambridge

Sadasivam, S. and Manicka, A. (2015). 'Biochemical Methods'. Third edition. New age international (p) limited, Publishers, New Delhi. 201- 202 pp.

Saha, S.; Mandal, A.; and Sahoo, D. (2017). Study of physico- chemical parameters of three different urban pond water of Nadia district, West- Bengal, India. *International Journal of Fisheries and Aquatic Studies*, 5(6): 23-27.

Sakhare,V. B. (2007). Reservoir Fisheries and Limnology, Narendra Publishing House, Delhi. 187pp.

Salaskar, P.B. (1998). Some environmental aspect of Powai Lake, Ph. D. Thesis, Mumbai University, Mumbai.

Sand, B.J.; Cheti, M.; Verma, V.K. and Kumar, D.(2012). Eco-hydrobiology with Special Amphasis on Ichthyofaunal Diversity of Urpod Wetland of Goalpara, Assam, India. *International Journal of Plant, Animal and Environmental Science*, 2(3): 103-109.

Santra, S.C.(2012). Environmental Science Published by New Central Book Agency (p) Ltd. 8/1 Chintamoni Das Lane, Kolkata, 1-638 pp.

Sarma, S.K.(2008). Wetlands: nature's precious gift. Published by The Assam Tribune (English daily), Guwahati, Assam, India, Saturday, February 2, Col.2, 6 p.

Sarma, D.; Das, J.; Goswami, U.C. and Dutta, A. (2012). Present Status and Habitat Ecology of *Ompok pabo* (Ham- Buchanan) in Goronga Beel, Morigaon, Assam (India). *Advances in Applied Science Research*, 3 (1): 481-488.

Sarma, S.K. and Borah,B.(2014). Phytosociological investigation of aquatic macrophytes of five wetlands of Sonitpur District of Assam, India. *Annals of Biological Research*, 5(9): 38-45.

Sarma, S.K. and Saikia, M. (2010). Utilization of wetland resources by the rural people of Nagaon district, Assam, *Indian Journal of Traditional Knowledge*, 9(1):145-151.

- Sarma, S.K. and Deka, U. (2014). Present status of aquatic macrophytes of the wetlands of Nalbari district of Assam, India. *Annals of Biological Research*, 5(5): 77- 84.
- Sarkar, U.K. and Ponniah, A.G. (2000). Evaluation of Northeast Indian Fishes for their potential. As Cultivable, Sport and Ornamental Fishes along with their Conservation and Endemic Status. 11-30p. In: Ponniah AG, Sarkar UK, editors. *Fish diversity of Northeastern India*. No-1. NBFGR, NATP publication Lucknow, U.P. 2-228 pp.
- Sarkar, S.; Deka, D. and Devi, N. (2008). Studies on some medicinally important wetland angiosperms used by the Bodo tribe of Kamrup District in Assam, India. *Pleione.*, 2(1): 20-26.
- Sartory, D.P. and Grobbelaar, J.U. (1984). Extraction of chlorophyll-a from fresh water phytoplankton for Spectrophotometric analysis. *Hydrobiologia*, 114:177-187.
- Sarwar, S.G. (1999). Water quality and periphytic algae component of Anchar Lake, Kashmir, Pages 237 – 250. In : Vijay Kumar, K. Freshwater ecosystem of India, Daya Publishing House, Delhi, 1- 336 pp.
- Saupi, N.; Zakira, M.H.; Bujang, J.S. (2009). Analytic Chemical composition and mineral content of yellow Velvetleaf (*Limnocharis flava* L. Buchenau) edible parts. *Journal of Applied Sciences*, 9: 2969-2974.
- Saupi, N.; Zakaria, M.H.; Bujang, J.S. and Arshad, A. (2015). The proximate compositions and mineral contents of *Neptunia oleracea* Loureiro, an aquatic plant from Malaysia. *Emirates Journal of Food and Agriculture*, 27(3): 266-274.
- Sculthorpe, C.D. (1967). The Biology of Aquatic Vascular Plants. Arnold, London. 610 pp.
- Seethal Lal, S.; Jaya, D.S. and Sherlywilliams, E. (2014). Ichthyofaunal diversity of Vattakkayal, a part of Ashamudi Lake, Kallam district, Kerala, South India. *Journal of Aquatic Biology and Fisheries*, 2: 620-626.

- Seethal, L. S.; Jaya, D.S. and Sherly Williams, E. (2013). Fish faunal Diversity of Vattakkayal, A part of Ashtamudi Lake, Kollam District, Kerala, South India. *International Journal of Scientific & Engineering Research*, 4(12), 418
- Sen, D.N. and Chatterjee, U.N. (1959). Ecological studies on aquatic and swampy vegetation of Gorakhpur. *Agra University Journal of Research (Science)*, 8(1): 17-29
- Sen, N. (2000). Occurrences, distribution and statues of diversified fish fauna of North East India. In: A.G. Ponniah and U.K. Sarkar (eds.) Fish Biodiversity of North East India, NBFGR, NATP Publication, 2-228 pp.
- Sen, N.(1982).Studies on the Systematics, Distribution and Ecology of the Ichthyofauna of Meghalaya and their bearing on the Fish and Fisheries of the State, vi +576,Ph.D. Thesis, Department of Zoology ,University of Gauhati (Assam)
- Sen, T.K. (1985). The fish fauna of Assam and the neighboring North Eastern States of India. Occasional Paper No. 64, Records of Zoological Survey of India. 1-216 pp.
- Setiawan, L. (1996). Effect of *Tribulus terrestris* L. on sperm morphology. Airlangga University, Surabaya, Indonesia
- Shahid, A.; Wani, K-Ul-Yaqoob and Pandit,A. K.(2007).Diversity and Distribution of Emergent Macrophytes in Nilnag Lake, a pine Forest Lake in Western Himalaya. *Journal of Research & Development*, 7:45-50.
- Shaikh, S. D. and Dongare, M. (2008). Analysis of photosynthesis pigments in Adiantum lunulatum Burm. At different localities of Sindhudurg District (Maharastra), *Indian Fern Journal*, 25: 83-86.
- Sharma, P. (1993). Geo-Ecological study of Beels and Swamps in Nagaon and Morigaon Districts, Assam. Ph. D. Thesis. Gauhati University, Assam.

- Sharma, A. and Singhal, P.K. (1998). Impact of floating and emergent vegetation on the trophy status of a tropical lake. The macrophytes and physic-chemical status. *Journal of Environmental Biology*, 9(3 suppl.): 303-311.
- Sharma, B.K. (2004). Phytoplankton Communities of a floodplain lake of the Brahmaputra River basin Upper Assam. *Journal of Indian Fisheries Association*, 31: 27-35
- Shaw, G.E. and Shebbeare, E.O. (1937). The fishes of northern Bengal. *Journal Royal Asiatic Society of Bengal, Science*, 5(3): 35-36.
- Shivaraju, M.; Warlu, V. and Ahmad, S. (2018). Fish diversity in Durgadahalli Lake of Tumakuru, Karnataka, India. *International Journal of Academic Research and Development*, 3(1): 1128-1131.
- Shraddha, S.; Rakesh, V.; Savita, D. and Praveen, J. (2011). Evaluation of water quality of Narmada River with reference to Physico-chemical parameters at Hoshangabad city, M.P. India. *Research Journal of Chemical Science*, 1(3).
- Shivananda, M. and Devaraj, K.V. (1991). Comparison of growth of carps fed on *Salvinia* based feed and conventional feed. *Fish Technology*, 28: 106-110.
- Shrivastava, N.P. and Bhattacharjya, B.K. (2003). Status of fisheries of flood plain wetlands of Assam, Pp. 22-29. In: Fisheries management of floodplain wetlands in India (Vinci, G.K., Jha, B.C.
- Shukla, R.S. and Chandel, P.S.(2014). A Textbook of plant Ecology, Published by S. Chand & Company Pvt. Ltd., New Delhi, 526 pp.
- Silva, S.S.; De.and Gunasekera, R.M. (1991).An evaluation of the growth of Indian and Chinese major carps in relation to the dietary protein content. *Aquaculture*, 92; 237-241.
- Singh, A.K. (2006). A contribution to the Aquatic and wetland flora of Varanasi. *Journal of Economic and Taxonomic Botany*, 30(1): 6-24.

- Singh, A.K. and Jayakumar, S. (2016). Water Quality Assessment of Kanwar Lake (Largest natural oxbow lake in the Indian subcontinent), Begusarai, Bihar, India. *Imperial Journal of Interdisciplinary Research (IJIR)*, 2(4): 793-803.
- Singh, A.K.; Tiwari, R.K.; Kanaujia, D.R. and Mishra, P. (2009). Physico-Chemical Characteristics of Ganga river water at Varanasi. *Journal of Ecobiology*, 25(I): 45- 46.
- Singh, B. K. and Jha, N.N. (2000)." Monthly variation in physic- chemical properties of sediment of Karwar Lake of North Bihar, India". *Indian Journal Environ and Ecoplan* 3 (3):725-728.
- Singh, K.K.; Sharma, B.M. and Usha, K.H. (2010). Ecology of Kharungpat lake, Thiribal, Manipur, India: Part 1 water quality status. *The Ecoscan*, 4(2 & 3): 241- 245.
- Singha, N.; Nag, R. and Deka, P. (2017). A Preliminary Study on Ichthyofaunal Diversity of Diplai Beel of Kokrajhar District of Assam, India. *International Journal of Fisheries and Aquatic Studies*, 5(3): 269-275.
- Sinha, S.N. and Biswas, M. (2011). Analysis of physic- chemical characteristics to study the water quality of a lake in Kalyani, West Bengal, *Asian Journal of Experimental Biological Sciences*, 2 (1): 18- 22.
- Singhal, R.N.; Jeet, S.; Davies, R.W.(1986). The physico-chemical environment and the plankton managed ponds in Haryana, India. Proceedings of the Indian Academy of Sciences (Animal Sciences), 95(3):353.
- Sivani, G.; Reddy, D.C. and Bhaskar, M. (2013). Effect on *Nymphaea* meal incorporated diets on growth feed efficiency and body composition in fingerlings of *Cyprinus carpio* L. *Journal of Applied and Natural Sciences*, 5(1): 5-9.
- Smith, P.F. (1962). Mineral analysis of plant tissues. *Annual Review of Plant Physiology*, 13: 81-108.

- Sneed, K.E.; Hastings, W. H. and Dupree, H.K.(1972). 'Warm water fish nutrition and future priorities'. *Resour. Publ. U.S. Bull. Sport Fish Wild.* 102: 135-140.
- Steffens, W. (1989). Principles of Fish Nutrition. Ellis Horwood, Chichester, 1-384 pp..
- Subramanyam, K. (1962). Aquatic angiosperms. Botanical Monograph (3), CSIR, New Delhi
- Sugunan, V.V. (1997). Fisheries management of small bodies in seven countries in Africa, Asia and Latin America. *FAO Circular*, No. 933:1- 149 pp.
- Sugunan,V.V. and Bhattacharjya, B.K.(2000). Ecology and Fisheries of Beels in Assam. Bull No 104, Cent. Inland Fish Res. Inst., Barrackpore, 65 pp.
- Suresh,V. (2003).Tilapias in: Lucas, J.S. and Southgate, P.C.,(Eds.) Aquaculture: farming of aquatic animals and plants, United Kingdom; Blackwell Publishing
- Sushil, M.; Reshi, J.M. and Krishna, M.(2014).Water quality status and responsible factors for variation in Anchart lake, Kashmir. *Journal of Environmental Science, Toxicology and FoodTechnology*, 8(2): 55-62.
- Steffens, W. (1981). Protein utilization of rainbow trout (*Salmo gairdneri*) and carp (*Cyprinus carpio*): A brief review, *Aquaculture*, 23: 337- 345.
- Swarnalatha, N. and Narasingarao, A.(1998).'Ecology studies of Banjara Lake with reference to water pollution'. *Journal of Environmental Biology*, 19(2): 179-186.
- Syed, M.A. and Coombs, T.L. (1982). Copper metabolism in the plaice *Pleuronectes platessa* (L). *Journal of Experimental Marine Biology and Ecology*, 63: 281- 286.
- Tacon, A.G.J. (1987).The nutrition and feeding of farmed fish and shrimp- A training Manual 1. The essential nutrients. A report prepared for the FAO trust fund GCP/RLA/O75/ITA project. Food and Agriculture organization of the United Nations, Brasilia, Brazil.

- Tacon, A. G. J. (1990). Standard methods for the nutrition and feeding of farmed fish and shrimp. Washington DC, Argent Laboratories press
- Talwar, P.K. and Jhingram, A.G. (1991). Inland Fishes of India and adjacent Countries, vol.1&2, Oxford and IBH Publishing House, Co. Pvt. Ltd., New Delhi, 1097 pp.
- Tamuli, P.; Bhagabati, S.K.; Dutta, R.; Zaman, ASN.; Sarma, J.; Choudhury, A.; Mudoi, L.P. and Debnath, D.J. (2018). Physico-chemical, characteristics of 47 No. Morakolong Beel, Morigaon district, Central Brahmaputra Valley Zone, Assam. *International Journal of chemical studies*, 6(5): 748-754.
- Tan, Y.T. (1970). Composition and nutritive value of some grasses, plants and aquatic weeds tested as diets. *Journal of Fish Biology*, 2: 253-257.
- Taylor, K.G. and Robbins, R.C. (1968). The amino acids composition of water hyacinth (*Eichhornia crassipes*) and its value as a protein supplement. *Hyacinth Cont. Journal*, 7: 24-25.
- Tripathi, A.K. and Gautam, M. (2007). Biochemical parameters of plants as indicators of air pollution. *Journal of Environmental Biology*, 28: 127-132.
- Tripathi, S.K. (1985). Hydrographical studies in Chilka Lake a Tropical lagoon. M. Phil. Dissertation, Berhampur University, Orrisa.
- Turan, M.; Kordali, S.; Zengin, H.; Dursun, A. and Sezen, Y. (2003). Macro and micro mineral content of some wild edible leaves consumed in Eastern Anatolia. *Acta Agriculturae Scandinavica, Section B, Plant Soil Science*, 53: 129-137.
- Turner, R.K.; Vander Bergh, J.C.J.M.; Soderqvist, T.; Barendregt, A.; Van der Straaten, J.; Maltby, E.; Van Ierland, E.C. (2000). Ecological-economic analysis of wetlands: Scientific integration for management and policy. *Ecological Economics*, 35 (1): 7-23.
- Umar, K.J.; Hassan, L.G.; Dangoggo, S.M. and Ladan, M.J. (2007). Nutritional Composition of Water Spinach (*Ipomoea aquatica* Forsk.) Leaves. *Journal of Applied Sciences*, 7 (6): 803-809.

United Nations Educational, Scientific and Cultural Organization(UNESCO, 2007). Case Studies on climate change and World Heritage. UNESCO World Heritage Centre, France.

Unni, K.S.(1971).An ecological study of the macrophytic vegetation of Doodhadhari Lake, Rajpur, M.P. India 1.Distribution and seasonal changes in aquatic plants. *Hydrobiologia*, 37:139-155.

Unni, K.S. (1967).Studies on the vegetation of ponds, swamps and river banks in Rajpur, M.P. *Journal of Bombay Natural History Society*, 64: 95-102.

United Nations Educational, Scientific and Cultural Organization (UNESCO, 2007).Case Studies on climate change and World Heritage. UNESWorld Heritage Centre, France.

USDA (2000). *Ipomoea aquatic* Forssk.United States Department of Agriculture. [www.ars-grin.gov/cgi-bin/npgs/htm/tax\\_search.pl?Ipomoea](http://www.ars-grin.gov/cgi-bin/npgs/htm/tax_search.pl?Ipomoea). Retrieve on 30th August, 2005.

Varghese, T.J.; Devaraj, K.V.; Shantharams and Shetty, H.P.C. (1976). ‘Growth response of the common Carp (*Cyprinus carpio* var. Com) to protein rich pelleted feed’ Symp. *Development and Utilization of Inland Fishery Resources*. Colombo (Sri Lanka), 408- 416 pp.

Vas, K.; Goud, J.V.; Suryam, A. and Singara Charya, M.A. (2009). Biomolecular and phytochemical analyses of three aquatic angiosperms. *African journal of Microbiology Research*, 3(8): 418-421.

Van Dijk, J.P. (1975). Iron metabolism in the tench (*Tinca tinca* L.). I. studies by means of intravascular administration of <sup>59</sup> Fe (III) bound to plasma. *Journal of Comparative Physiology*, 99: 321- 330.

Vasu, K.; Jakku,V. G.; Suryam, A.;Charya, S. M. (2009). Bimolecular and photochemical analysis of three aquatic angiosperm. *African Journal of Microbiology Research*, 3(8):418-421.

- Venkateshwarlu, M.; Somashekar, D.S. and Shanmugam, M.(2002)."Influence of Hydrophyte leaf-based feed on the growth of Carp fry". Wetlands Conservation and Management. Published by Pointer Publishers, Jaipur (Raj), India.
- Venugopal, M.N.and Keshavanath, P. (1984)."Influence of supplementary feeds on the biochemical composition on the freshwater carps, *Catla catla* (Ham.), *Cirrhinus mrigala* (Ham.) and *Cyprinus carpio* (Ham.)". *Indian Journal of Animal Sciences*, 54(6): 555-559.
- Verma, D.M. (1971). Cyperaceae of Assam and Neighbouring areas; Ph.D. Thesis. Gauhati University, Guwahati, Assam.
- Vicas, S.I.; Laslo, V.; Pantea, S. and Bandict, G.E. (2010). Chlorophyll and carotenoids pigments from Mistletoe (*Viscum album*) leaves using different solvents. *Fascicula Biology*, xvii (2): 213-218.
- Vijayalakshmi, G.and Sivapragasam,C.(2008).Research Methods Tips and Techniques, MJP Publishers, Chennai, 218 pp.
- Vishwanath, W.; Manojkumar, W.; Kosygin L. and Selim, K.S. (1998). Biodiversity of freshwater fishes of Manipur, India. *Italian Journal of Zoology*, 65 (1): 321-324.
- Vishwanath,W.; Lakra,W.S.and Sarkar,U.K. (2007)."Fishes of North East India". Publisher National Bureau of Fish Genetic Resource, Indian Council of Agricultural Research.
- Vishwanath, W.; Ng, H.H.; Britz, R.; Singh,L.K.; Chaudhury,S. and Conway, K.W.(2010). In: D.J. Allen, S. Molur; B.A.Daniel(Ed), The status and distribution of freshwater biodiversity in the Eastern Himalaya, 22-39 pp.
- Vyas, L.N. (1964). A study of hydrophytes and marsh plants of Alwar and environs. *Journal of Indian Botanical Society*. 43: 7-30.
- Wani, O.A. and Gupta, U.S. (2015). A Study on Ichthyofaunal Diversity of Sagar Lake, Madhya Pradesh, India. *International Journal of Biodiversity and Conservation*. 7(3): 126-129.

Wani, S.A.; Yaqoob, K.U. and Pandit, A.K. (2007). Diversity and Distribution of Emergent Macrophytes in Nilnag Lake, a pine Forest lake in Western Himalaya, *Journal of Research and Development*, Vol. 7:45

Wasagu, R.S.U.; Lawal, M.; Galadima, L.G. and Aliero, A.A. (2015). Nutritional composition, antinutritional factors and elemental analysis of *Nymphaea lotus* (water lily), *BayeroJournal of Pure and Applied sciences*, 8(1): 1-5.

Watanabe, T.; Satoh, S. and Takeuchi, T. (1988). Availability of minerals in fishmeal to fish. *Asian Fisheries Society*, 1(2): 175- 195 pp.

Watanabe, T.; Kiron, V. and Satoh, S. (1997). Trace minerals in fish nutrition. *Aquaculture*, 151: 185-207.

Website : www. Fishbase. Org.

Weaver, J.E. and Clements, F.E. (1929). Plant Ecology, 1st ed. Published by Mc Graw-Hill Book Company, New York, 520 pp.

Wee, K.L. and Wang, S.S. (1987). Nutritive value of Leucaena leaf meal in pelleted feed for Nile Tilapia. *Aquaculture*, 62: 97-108.

Weir, T. and Stocking, C. (1952). The chloroplast: structure, inheritance and enzymology. *Botanical Review*, 18:14

Whigham, D.F.; Dykyjova, D.; Hejny, S. (1993). Wetlands of the world I: Inventory, ecology and management, Kluwer Academic Publishers, The Netherlands, eds.

Wetzel, R.G. (2001). Limnology: Lake and river ecosystems, 3<sup>rd</sup>ed. Academic press, Elsevier, San Diego. New York. 1006 pp.

WHO (1995). World Health organization, Guideline for Drinking water quality. Geneva

WHO (1971). International standard for Drinking water. II end Geneva; World Health Organization.

Wilson, N.M. (1995). Nutritional and feeding of fish, New York

- Wolken, J. (1961). *Euglena*: An Experimental Organism for Biochemical and Biophysical Studies, New Brunswick, N.J.: Rutgers University Press.
- Yadavi, P.; Yadavi, V.K.; Yadavi, A.K. and Khare, P.K. (2013). Physico-Chemical Characteristics of a Fresh Water Pond of Orai, U.P., Central India. *Octa Journal of Biosciences*, 1(2): 177-184.
- Zacharias, V.J.; Bhardwaj, A.K. and Jacob, P.C. (1996). Fish fauna of Periyar Tiger Reserve. *Journal of the Bombay Natural History Society*, 93: 38-43.
- Zafar, A.R. (1964). On the ecology of algae in certain fish ponds of Hyderabad, India; physico- chemical Complex. *Hydrobiologia*. 23(2): 179-195.
- Zafar, M.; Khan, M.A.; Ahmad, M.; Jain, G.; Sultan, S.; Ullah, K.; Marwat, S.K.; Ahmad, F.; Jabeen, A.; Nazir, A.; Abbasi, A.; Rehman, Z. and Ullah, Z. (2010). Elemental analysis of some medicinal plants used in traditional medicine by atomic absorption spectrophotometer (AAS). *Journal of Medicinal Plants Research*, 4(19): 1987-1990.
- Zutshi, D.P.; Subia, B.A.; Khan, M.A. and Wanganeo, A. (1980). "Comparative limnology of nine lakes of Jammu and Kashmir, Himalayas". *Hydrobiologia*, 72(1-2): 101-112.