# A STUDY ON PHARMACOLOGICAL ACTIVITIES, NUTRITIONAL VALUE AND DNA BARCODING OF SELECTED ETHNOMEDICINAL PLANTS FROM GOALPARA DISTRICT, ASSAM



A thesis submitted to Bodoland University for the degree of Doctor of Philosophy in Biotechnology in the Faculty of Science and Technology, 2024

BY

SEEMA KHAKHALARY Registration No: FINAL.BIO00212 of 2019-2020 Department of Biotechnology Bodoland University, Kokrajhar- 783370 Assam, India

## **DECLARATION**

I, Seema Khakhalary, do hereby declare that the thesis entitled "A Study On Pharmacological Activities, Nutritional Value and DNA Barcoding of Selected Ethno Medicinal Plants from Goalpara District, Assam," is the original research work done by me in design and execution, that all material contained herein has been duly acknowledged, and that the thesis hereby submitted to Bodoland University for the degree of Doctor of Philosophy in Biotechnology has not been previously by me for a degree at this or any other University.

Date: Place: Kokrajhar (SEEMA KHAKHALARY)

From, Dr. Silistina Narzari Department of Biotechnology Bodoland University Kokrajhar, Assam Email: nsilistina@gmail.com

**CERTIFICATE** 

This is to certify that the thesis work entitled, "A Study on Pharmacological Activities, Nutritional Value and DNA Barcoding of Selected Ethno Medicinal Plants from Goalpara District, Assam," submitted by Seema Khakhalary (FINAL.BIO00212 of 2019-2020) in partial fulfillment of the requirement for the degree of Doctor of Philosophy in Biotechnology is a record of bonafide work carried out by her under my supervision and guidance.

(SILISTINA NARZARI) Supervisor & Assistant Professor Department of Biotechnology Bodoland University

Dated:

### **ACKNOWLEDGEMENT**

First and foremost, I offer praises and thanksgivings to the Almighty, for his showers of blessings throughout my research work to complete the research successfully.

My heartfelt gratitude to **Prof. L. L. Singh** (Former), and **Prof. B. L Ahuja** (Present), **Vice Chancellor, Dr. S Basumatary**, Registrar, **Dr. M. Basumatary**, Academic Registrar, Bodoland University for giving me the opportunity and environment to initiate and develop my research work.

I owe my deepest gratitude to my supervisor Dr. Silistina Narzari, Assistant Professor, Department of Biotechnology, Bodoland University, Kokrajhar for her valuable guidance, suggestions, encouragement and insightful comments which were vital for my research work. It was a great privilege and honor to work and study under her guidance.

With respect and complacency I sincerely thank **Prof. Sandeep Das**, the Head and other faculty members. **Prof. Jatin Sarmah**, **Dr. Arvind Kumar Goyal**, **Dr. Sharmistha Brahma Kour**, **Mrs. Jonali Brahma** and **Dr. Monoranjan Boro** for their kind support and help in various way throughout this course.

I owe my hearty gratitude to Ministry of Tribal Affairs, Government of India for providing me fellowship under the scheme of National Fellowship for Schedule Tribe (NFST) to carry out my research work. I am also highly indebted to Assam Institute of Research for Tribals and Schedule Caste for granting me financial assistance for my research work.

I sincerely acknowledge and thank Biotech Park, IIT, Guwahati for providing laboratory facilities for doing GCMS and FTIR study. I am also thankful to Sophisticated Analytical Instrument Facility, NEHU, Shillong for helping me to perform ICP-OES mineral analysis. I would also like to offer my sincere thanks to Sandor Speciality Diagnostics Private Limited, Hyderabad for their support in UPLC analysis. I also express my acknowledgement to Mr. Biologist Private Limited, Pune for providing support in in-vitro anticancer study. My cordial thanks to the co-ordinators of the Advanced Level Institutional Biotech Hub and Bioinformatics Infrastructure Facility (BIF) of the Department of Biotechnology for assisting and providing laboratory facilities. Most importantly I would like to acknowledge the help from Dr. Saurabh Borah, Assistant Professor, Gauhati University Botanical Herbarium and Dr. Sanjib Baruah, Assistant Professor, Bodoland University Botanical Herbarium in taxonomic study of the plant for authentication. I farther take this opportunity to thank Dr. Tapan Dutta, Principal, Jawaharlal Nehru College, Boko for his various suggestions, and for offering the deepest knowledge and devotion whenever sought in this field during the tenure of the investigation.

Here I take this great opportunity to express my sincere thanks and gratitude to the **traditional healers**, local people and tribal inhabitants from the study sites for their valuable information, persistent assistance and guidance during my field study. I believe without the contributions form you all I would never have reached this far. I will always remain indebted towards you all for this.

My whole hearted thanks to my fellow researchers for their kind help, co-operation and for always being there to support me in all my good and bad days. The love and support made my journey towards completing this thesis easier.

Also, I sincerely thank Mr. Chandan Brahma, Mr. Hirakmoni Borgayary, and Mr. Paban Roy technical staff of Department of Biotechnology, Bodoland University, Kokrajhar for their constant support throughout this research work. With deepest affection, I attempt to extend my thanks to all the supporting staff of this institution for without their help the work would never have been completed.

Finally, I express my sincere thanks to all my family members for their strong support, perseverance and valuable prayers. I am extremely grateful to my parents for their love, prayers, caring and sacrifices for educating and preparing me for my future and continuing support to complete this research work. My special thanks goes to my dear father Mr. Binoy Ram Khakhalary and mother Mrs. Jamuna Khakhalary for the keen interest they showed to encourage me to complete this thesis successfully.

#### (SEEMA KHAKHALARY)

# LIST OF TABLES

Table 3.1	Primer used for PCR amplification and references used
Table 4.1	Medicinal plants used by the traditional healers of Goalpara, Assam
Table 4.2	Voucher specimen with accession number
Table 4.3	Identification of the plant species using BLASTn
Table 4.4	Proximate composition analysis of studied plants (g/100g) dry weight
Table 4.5	Recommended intake of essential minerals per day compared with <i>Z. oxyphyllum</i>
Table 4.6	Recommended intake of essential minerals per day compared with <i>R. serrata</i>
Table 4.7	Recommended intake of essential minerals per day compared with <i>B. lanceolaria</i>
Table 4.8	Amino acid profile of Z. oxyphyllum (in mg/100g)
Table 4.9	Amino acid score of <i>Z. oxyphyllum</i> based on FAO/WHO/UNU (1985) consultation pattern of requirement for a 2- 5 year preschool child
Table 4.10	Amino acid profile of <i>R. serrata</i> (in mg/100g)
Table 4.11	Amino acid score of <i>R. serrata</i> based on FAO/WHO/UNU (1985) consultation pattern of requirement for a 2- 5 year preschool child
Table 4.12	Amino acid profile of <i>B. lanceolaria</i> (in mg/100g)
Table 4.13	Amino acid score of <i>B. lanceolaria</i> based on FAO/WHO/UNU (1985) consultation pattern of requirement for a 2- 5 year preschool child

- Table 4.14Fatty acid identified in Z. oxyphyllum
- Table 4.15Fatty acid identified in R. serrata
- Table 4.16Fatty acid identified in B. lanceolaria
- Table 4.17Anti nutritional contents of plant species (in mg/100g dry weight)
- Table 4.18Yield percentage (%) of the plant species extracted using different<br/>solvents
- Table 4.19Phytochemicals present in Zanthoxylum oxyphyllum extracts
- Table 4.20Phytochemicals present in Rotheca serrata extracts
- Table 4.21
   Phytochemicals present in Blumea lanceolaria extracts
- Table 4.22Total phenolic content of plant extracts
- Table 4.23Total flavonoid content of plant extracts
- Table 4.24 IC50 values (µg/mL) of *Z. oxyphyllum*, *R. serrata* and *B. lanceolaria* in DPPH assay
- Table 4.25 IC50 values ( $\mu$ g/mL) of *Z. oxyphyllum*, *R. serrata* and *B. lanceolaria* in H<sub>2</sub>O<sub>2</sub> assay
- Table 4.26Antioxidant activity of Z. oxyphyllum, R. serrata and B. lanceolariain Ferric Reducing Power Assay
- Table 4.27
   Bioactive compounds identified in aqueous extract of Z. oxyphyllum
- Table 4.28Bioactive compounds identified in hexane extract of Z. oxyphyllum

#### Table 4.29 Bioactive compounds identified in chloroform extract of Z. oxyphyllum

- Table 4.30
   Bioactive compounds identified in methanol extract of Z. oxyphyllum
- Table 4.31
   Bioactive compounds identified in aqueous extract of R. serrata
- Table 4.32
   Bioactive compounds identified in hexane extract of R. serrata

- Table 4.33
   Bioactive compounds identified in chloroform extract of R. serrata
- Table 4.34 Bioactive compounds identified in methanol extract of *R. serrata*
- Table 4.35
   Bioactive compounds identified in aqueous extract of B. lanceolaria
- Table 4.36
   Bioactive compounds identified in hexane extract of B. lanceolari
- Table 4.37 Bioactive compounds identified in chloroform extract of B.
   Ianceolaria
- Table 4.38
   Bioactive compounds identified in methanol extract of *B. lanceolaria*
- Table 4.39
   FTIR peak values and functional groups in Zanthoxylum oxyphyllum leaf
- Table 4.40 FTIR peak values and functional groups in Rotheca serrata leaf
- Table 4.41 FTIR peak values and functional groups in *Blumea lanceolaria* leaf
- Table 4.42Cytotoxicity activity of the plant extracts against MCF-7 and HeLa<br/>cell lines
- Table 4.43
   Quantitative apoptotic analysis against MCF-7 Cells using AO/Et-Br dual staining
- Table 4.44 Quantitative apoptotic analysis against HeLa Cells using AO/Et-Br dual staining
- Table 4.45 Quantitative cell distribution of MCF-7 cells over the treatment of the plant extracts in various apoptotic stages
- Table 4.46Quantitative cell distribution of HeLa cells over the treatment of B.lanceolaria in various apoptotic stages

## **LIST OF FIGURES**

- Figure 4.1 Maximum-Likelihood phylogenetic tree of similarity between *Z. oxyphyllum* and the closest species generated by Kimura 2 parameter using rbcL gene
- Figure 4.2 Maximum-Likelihood phylogenetic tree of similarity between *R*. *serrata* and the closest species generated by Kimura 2 parameter using ITS1 gene
- Figure 4.3 Maximum-Likelihood phylogenetic tree of similarity between *B. lanceolaria* and the closest species generated by Kimura 2 parameter using ITS2 gene
- Figure 4.4 UPLC chromatogram of amino acid (Standard)
- Figure 4.5 UPLC chromatogram of amino acid in Z. oxyphyllum
- Figure 4.6 UPLC chromatogram of amino acid in *R. serrata*
- Figure 4.7 UPLC chromatogram of amino acid in *B. lanceolaria*
- Figure 4.8 GCMS chromatogram of fatty acids in Z. oxyphyllum
- Figure 4.9 GCMS chromatogram of fatty acids in *R. serrata*
- Figure 4.10 GCMS chromatogram of fatty acids in *B. lanceolaria*
- Figure 4.11 Standard gallic acid calibration curve for total phenolic content (TPC)
- Figure 4.12 Standard quercitin calibration curve for total flavonoid content (TFC)
- Figure 4.13 Standard graph (Ascorbic acid) in Ferric Reducing Power Assay
- Figure 4.14 GC-MS chromatogram of aqueous extract of Z. oxyphyllum
- Figure 4.15 GC-MS chromatogram of hexane extract of Z. oxyphyllum

- Figure 4.16 GC-MS chromatogram of chloroform extract of Z. oxyphyllum
- Figure 4.17 GC-MS chromatogram of methanol extract of Z. oxyphyllum
- Figure 4.18 GC-MS chromatogram of aqueous extract of *R. serrata*
- Figure 4.19 GC-MS chromatogram of hexane extract of *R. serrata*
- Figure 4.20 GC-MS chromatogram of chloroform extract of *R. serrata*
- Figure 4.21 GC-MS chromatogram of methanol extract of *R. serrata*
- Figure 4.22 GC-MS chromatogram of aqueous extract of *B. lanceolaria*
- Figure 4.23 GC-MS chromatogram of hexane extract of *B. lanceolaria*
- Figure 4.24 GC-MS chromatogram of chloroform extract of *B. lanceolaria*
- Figure 4.25 GC-MS chromatogram of methanol extract of *B. lanceolaria*
- Figure 4.26 FTIR spectrum of Z. oxyphyllum leaves
- Figure 4.27 FTIR spectrum of *R. serrata* leaves
- Figure 4.28 FTIR spectrum of *B. lanceolaria* leaves
- Figure 4.29 % of inhibition against MCF-7 cell line
- Figure 4.30 % Cell viability of promising extracts against MCF-7 cancer cells
- Figure 4.31 % inhibition against HeLa cell line
- Figure 4.32 % Cell viability of promising extracts against HeLa cancer cells

### LIST OF IMAGES

- Plate 3.1 Map of Goalpara showing study area
- Plate 4.1 Zanthoxylum oxyphyllum Edgew
- Plate 4.2 *Rotheca serrata* (L.) Steane & Mabb.
- Plate 4.3 *Blumea lanceolaria* (Roxb.) Druce
- Plate 4.4 1.5% agarose gels showing amplified products of rbcL (A) for Z. oxyphyllum, ITS1 (B) for R. serrata and ITS2 (C) for B. lanceolaria
- Plate 4.5 Alignment view of Z. oxyphyllum using combination of NCBI GenBank
- Plate 4.6 Sequences of Z. oxyphyllum showing significant alignments
- Plate 4.7 Alignment view of *R. serrata* using combination of NCBI GenBank
- Plate 4.8 Sequences of *R. serrata* showing significant alignments
- Plate 4.9 Alignment view of *B. lanceolaria* using combination of NCBI GenBank
- Plate 4.10 Sequences of *B. lanceolaria* showing significant alignments
- Plate 4.11 Morphological changes of MCF-7 cells over treatment of the plant extracts
- Plate 4.12 Morphological changes of HeLa cells over treatment of the plant extracts
- Plate 4.13 Apoptotic Cell death analysis of MCF-7 cells over the treatment of Met\_SKO-1(*Z.oxyphyllum*), Met\_SKO-2 (*R. serrata*) and Met\_SKO-3 (*B. lanceolaria.*) at their IC<sub>50</sub> and 2xIC<sub>50</sub> using AO/Et-Br staining method. Doxorubicin was used as a positive control
- Plate 4.14 Apoptotic Cell death analysis of HeLa cells over the treatment of Met\_SKO-1(*Z.oxyphyllum*), Met\_SKO-2 (*R. serrata*) and Met\_SKO-3 (*B. lanceolaria.*) at their IC<sub>50</sub> and 2xIC<sub>50</sub> using AO/Et-Br staining method. Doxorubicin was used as a positive control
- Plate 4.15 Flow cytometry result of the apoptotic activities of the samples against MCF-7 cancer cell
- Plate 4.16 Flow cytometry result of the apoptotic activity of *B. lanceolaria* against HeLa cancer cell

## **LIST OF ABBREVIATIONS**

AIDS	-	Acquired Immune Deficiency Syndrome
AO/Et-Br	-	Acridine orange/Ethidium bromide
AOAC	-	Association of Official Analytical Chemists
BC	-	Before Christ
BSI	-	Botanical Society of India
BLAST	-	Basic Local Alignment Search Tool
CTAB	-	Cetyl Trimethyl Ammonium Bromide
CTM	-	Chinese Traditional Medicine
CuSO <sub>4</sub>	-	Copper Sulphate
dNTP	-	Deoxynucleoside triphosphates
DMSO	-	Dimethyl sulphoxide
DNA	-	Deoxyribonucleic acid
EAA	-	Essential Amino Acid
EDTA	-	Ethylenediamine tetra acetic acid
FACS	-	Flourescence activated cell sorting
FAO	-	Food and Agriculture Organization
FBS	-	Fetal Bovine Serum
V-FITC	-	Fluorescein Isothiocyanate
FRAP	-	Ferric Reducing Antioxidant Power Assay
FT-IR	-	Fourier Transformed Infrared Spectroscopy
GA	-	Gallic Acid
GCMS	-	Gas Chromatography Mass Spectrometry
HCl	-	Hydrochloric Acid
HIV	-	Human Immuno Deficiency Virus
$H_2SO_4$	-	Sulfuric Acid
$H_2O_2$	-	Hydrogen Peroxide
HNO <sub>3</sub>	-	Nitric Acid
IC50	-	Inhibitory Concentration
ICP-OES	-	Inductively Coupled Plasma-Optical Emission Spectrometry

ISSR	-	Inter-simple sequence repeats
ITS	-	Internal transcribed spacer
$K_2SO_4$	-	Potassium Sulphate
KMnO <sub>4</sub>	-	Potassium Permanganate
М	-	Molar
MEGA	-	Molecular Evolutionary Genetics
MTT	-	3-[4, 5-dimethylthiazol-2-yl]-2, 5 diphenyl tetrazolium bromide)
Ng	-	nanogram
NaCl	-	Sodium Chloride
NaOH	-	Sodium Hydroxide
NCBI	-	National Centre for Biotechnology Information
NEAA	-	Non-Essential Amino Acid
NER	-	Northeastern Region
$N_2$	-	Nitrogen
NIST	-	National Institute of Standard and Technology
PCR	-	Polymerase Chain Reaction
PBS	-	Phosphate Buffered Saline
PI	-	Propidium Iodide
RAPD	-	Random Amplified Polymorphic DNA
rbcL	-	Ribulose-bisphosphate carboxylase
RBC	-	Red Blood Corpucles
RDA	-	Recommended dietary allowances
SD	-	Standard Deviation
TCM	-	Traditional Chinese Medicine
TFC	-	Total flavonoid content
UV-VIS	-	Ultra Violet Visible
WBC	-	White Blood Corpuscles
WHO	-	World Health Organisation