

# Bibliography

Abbott, T. M. C., Allam, S., Andersen, P., Angus, C., Asorey, J., Avelino, A., ... & DES Collaboration. (2019). First cosmology results using type Ia supernovae from the dark energy survey: constraints on cosmological parameters. *The Astrophysical Journal Letters*, **872(2)**, L30.

Abdel-Megied, M., & Hegazy, E. A. (2016). Bianchi type-VI cosmological model with electromagnetic field in Lyra geometry. *Canadian Journal of Physics*, **94(10)**, 992-1000.

Adhav, K. S. (2011). LRS Bianchi type-I cosmological model with linearly varying deceleration parameter. *The European Physical Journal Plus*, **126(12)**, 1-5.

Adhav, K. S., Bansod, A. S., Munde, S. L., & Nakwal, R. G. (2011). Bianchi type- $VI_0$  cosmological models with anisotropic dark energy. *Astrophysics and Space Science*, **332(2)**, 497-502.

Adhav, K. S., Dawande, M. V., Thakare, R. S., & Raut, R. B. (2011). Bianchi type-III magnetized wet dark fluid cosmological model in general relativity. *International Journal of Theoretical Physics*, **50(2)**, 339-348.

Ahmed, N., & Pradhan, A. (2014). Bianchi type-V cosmology in  $f(R, T)$  gravity with  $\Lambda(T)$ . *International Journal of Theoretical Physics*, **53(1)**, 289-306.

- Akarsu, Ö., & Dereli, T. (2012). Cosmological models with linearly varying deceleration parameter. *International Journal of Theoretical Physics*, **51(2)**, 612-621.
- Akarsu, Ö., & Kılınç, C. B. (2010). Bianchi type-III models with anisotropic dark energy. *General Relativity and Gravitation*, **42(4)**, 763-775.
- Aktaş, C., Aygün, S., & Sahoo, P. K. (2018). Relationship between magnetic field and anisotropy parameter in gravitation theories. *Modern Physics Letters A*, **33(24)**, 1850135.
- Amirhashchi, H., Zainuddin, H., & Pradhan, A. (2011). Magnetized Bianchi Type-III String Universe with Time Decaying Vacuum Energy Density  $\Lambda$ . *International Journal of Theoretical Physics*, **50(8)**, 2531-2545.
- Amirhashchi, H., Pradhan, A., & Jaiswal, R. (2013). Two-fluid dark energy models in Bianchi type-III universe with variable deceleration parameter. *International Journal of Theoretical Physics*, **52(8)**, 2735-2752.
- Amirhashchi, H. (2014). Interacting viscous dark energy in a Bianchi type-III universe. *Research in Astronomy and Astrophysics*, **14(9)**, 1121.
- Arkani-Hamed, N., Cheng, H. C., Luty, M. A., & Mukohyama, S. (2004). Ghost condensation and a consistent infrared modification of gravity. *Journal of High Energy Physics*, **2004(05)**, 074.
- Arkani-Hamed, N., Finkbeiner, D. P., Slatyer, T. R., & Weiner, N. (2009). A theory of dark matter. *Physical Review D*, **79(1)**, 015014.
- Armendariz-Picon, C., Mukhanov, V., & Steinhardt, P. J. (2000). Dynamical solution to the problem of a small cosmological constant and late-time cosmic acceleration. *Physical Review Letters*, **85(21)**, 4438.

Armendariz-Picon, C., Mukhanov, V., & Steinhardt, P. J. (2001). Essentials of k-essence. *Physical Review D*, **63(10)**, 103510.

Asgar, A., & Ansari, M. (2014). Accelerating Bianchi type- $VI_0$  bulk viscous cosmological models in Lyra geometry. *Journal of Theoretical and Applied Physics*, **8(4)**, 219-224.

Avelino, A., & Nucamendi, U. (2009). Can a matter-dominated model with constant bulk viscosity drive the accelerated expansion of the universe?. *Journal of Cosmology and Astroparticle Physics*, **2009(04)**, 006.

Bali, R., & Dave, S. (2001). Bianchi type IX string cosmological model in general relativity. *Pramana*, **56(4)**, 513-518.

Bali, R., & Dave, S. (2002). Bianchi type-III string cosmological model with bulk viscous fluid in general relativity. *Astrophysics and space science*, **282(2)**, 461-466.

Bali, R., & Dave, S. (2003). Bianchi type-IX string cosmological models with bulk viscous fluid in general relativity. *Astrophysics and space science*, **288(4)**, 399-405.

Bali, R., & Yadav, M. K. (2005). Bianchi Type-IX viscous fluid cosmological model in general relativity. *Pramana*, **64(2)**, 187-196.

Bali, R., & Pradhan, A. (2007). Bianchi type-III string cosmological models with time dependent bulk viscosity. *chinese physics letters*, **24(2)**, 585.

Bali, R., & Jain, S. (2007). Bianchi type-III non-static magnetized cosmological model for perfect fluid distribution in general relativity. *Astrophysics and Space Science*, **311(4)**, 401-406.

Bali, R., Pradhan, A., & Amirhashchi, H. (2008). Bianchi type- $VI_0$  magnetized barotropic bulk viscous fluid massive string universe in general relativity. *International Journal of Theoretical Physics*, **47(10)**, 2594-2604.

- Bali, R., Banerjee, R., & Banerjee, S. K. (2008). Bianchi type- $VI_0$  magnetized bulk viscous massive string cosmological model in General Relativity. *Astrophysics and space science*, **317(1)**, 21-26.
- Bali, R., Banerjee, R., & Banerjee, S. K. (2009). Some LRS Bianchi Type- $VI_0$  Cosmological Models with Special Free Gravitational Fields. *Electronic Journal of Theoretical Physics*, **6(21)**.
- Bali, R., Tinker, S., & Singh, P. (2010). Bianchi type III cosmological models for barotropic perfect fluid distribution with variable G and  $\Lambda$ . *International Journal of Theoretical Physics*, **49(7)**, 1431-1438.
- Bamba, K., Capozziello, S., Nojiri, S. I., & Odintsov, S. D. (2012). Dark energy cosmology: the equivalent description via different theoretical models and cosmography tests. *Astrophysics and Space Science*, **342(1)**, 155-228.
- Banerjee, N., & Das, S. (2005). Acceleration of the universe with a simple trigonometric potential. *General Relativity and Gravitation*, **37(10)**, 1695-1703.
- Baro, J., Singh, K. P., & Singh, A. T. (2021). Mathematical analysis on anisotropic Bianchi type-III inflationary string cosmological models in Lyra geometry. *Indian Journal of Science and Technology*, **14(1)**, 46-54.
- Barrow, J. D. (1984). Helium formation in cosmologies with anisotropic curvature. *Monthly Notices of the Royal Astronomical Society*, **211(2)**, 221-227.
- Barrow, J. D. (1990). Graduated inflationary universes. *Physics Letters B*, **235(1-2)**, 40-43.
- Behera, D., Tripathy, S. K., & Routray, T. R. (2010). Bulk Viscous Bianchi-III Models with Time Dependent G and  $\Lambda$ . *International Journal of Theoretical Physics*, **49(10)**, 2569-2581.

- Bento, M. C., Bertolami, O., & Sen, A. A. (2002). Generalized Chaplygin gas, accelerated expansion, and dark-energy-matter unification. *Physical Review D*, **66(4)**, 043507.
- Berman, M. S. (1983). A special law of variation for Hubble's parameter. *Nuovo Cimento B*, **74(2)**, 182-186.
- Berman, M. S., & de Mello Gomide, F. (1988). Cosmological models with constant deceleration parameter. *General Relativity and Gravitation*, **20(2)**, 191-198.
- Bernardeau, F., Colombi, S., Gaztanaga, E., & Scoccimarro, R. (2002). Large-scale structure of the Universe and cosmological perturbation theory. *Physics reports*, **367(1-3)**, 1-248.
- Bertone, G., Hooper, D., & Silk, J. (2005). Particle dark matter: Evidence, candidates and constraints. *Physics reports*, **405(5-6)**, 279-390.
- Bianchi, L. (1918). Lezioni sulla teoria dei gruppi continui finiti di trasformazioni. *E. Spoerri*, 550
- Bianchi, L. (1898). On the three-dimensional spaces which admit a continuous group of motions. *Memorie di Matematica e di Fisica della Società Italiana delle Scienze*, **11**, 267-352.
- Blandford, R. D., Amin, M., Baltz, E. A., Mandel, K., & Marshall, P. J. (2005). Cosmokinetics. *In Observing Dark Energy*. **339**, 27.
- Brans, C., & Dicke, R. H. (1961). Mach's principle and a relativistic theory of gravitation. *Physical review*, **124(3)**, 925.
- Bull, P., Akrami, Y., Adamek, J., Baker, T., Bellini, E., Jimenez, J. B., ... & Winther, H. A. (2016). Beyond  $\Lambda$ CDM: Problems, solutions, and the road ahead. *Physics of the Dark Universe*, **12**, 56-99.

- Burd, A. B., Buric, N., & Ellis, G. F. R. (1990). A numerical analysis of chaotic behaviour in Bianchi IX models. *General relativity and gravitation*, **22(3)**, 349-363.
- Caldwell, R. R. (2002). A phantom menace? Cosmological consequences of a dark energy component with super-negative equation of state. *Physics Letters B*, **545(1-2)**, 23-29.
- Caldwell, R. R., Komp, W., Parker, L., & Vanzella, D. A. (2006). Sudden gravitational transition. *Physical Review D*, **73(2)**, 023513.
- Carroll, S. M. (2001). The cosmological constant. *Living reviews in relativity*, **4(1)**, 1-56.
- Carroll, S. M., Hoffman, M., & Trodden, M. (2003). Can the dark energy equation-of-state parameter  $w$  be less than  $-1$ ?. *Physical Review D*, **68(2)**, 023509.
- Chakraborty, S. (1991). String cosmology in Bianchi- $VI_0$  space-time. *Indian Journal of Pure and Applied Physics*, **29(1)**, 31-33.
- Chand, A., Mishra, R. K., & Pradhan, A. (2016). FRW cosmological models in Brans-Dicke theory of gravity with variable  $q$  and dynamical  $\Lambda$ -term. *Astrophysics and Space Science*, **361(2)**, 1-12.
- Collins, C. B., Glass, E. N., & Wilkinson, D. A. (1980). Exact spatially homogeneous cosmologies. *General Relativity and Gravitation*, **12(10)**, 805-823.
- Collins, C. B., & Hawking, S. W. (1973). Why is the universe isotropic?. *The Astrophysical Journal*, **180**, 317-334.
- Copeland, E. J., Sami, M., & Tsujikawa, S. (2006). Dynamics of dark energy. *International Journal of Modern Physics D*, **15(11)**, 1753-1935.
- Cunha, J. V. (2009). Kinematic constraints to the transition redshift from supernovae type Ia union data. *Physical Review D*, **79(4)**, 047301.

- Cyburtt, R. H., Fields, B. D., Olive, K. A., & Yeh, T. H. (2016). Big bang nucleosynthesis: Present status. *Reviews of Modern Physics*, **88**(1), 015004.
- Debnath, U., Banerjee, A., & Chakraborty, S. (2004). Role of modified Chaplygin gas in accelerated universe. *Classical and Quantum Gravity*, **21**(23), 5609.
- De Oliveira, H. P., de Almeida, A. O., Soares, I. D., & Tonini, E. V. (2002). Homoclinic chaos in the dynamics of a general Bianchi type-IX model. *Physical Review D*, **65**(8), 083511.
- Dewri, M. (2018). Magnetized Anisotropic Bianchi Type-VI Cosmological Model Containing Dark Energy. *IOSR Journal of Applied Physics*, **10**(1), 31-35
- Dixit, A., Zia, R., & Pradhan, A. (2020). Anisotropic bulk viscous string cosmological models of the Universe under a time-dependent deceleration parameter. *Pramana*, **94**(1), 1-11.
- Dubinski, J., & Carlberg, R. G. (1991). The structure of cold dark matter halos. *The Astrophysical Journal*, **378**, 496-503.
- Dunn, K. A., & Tupper, B. O. J. (1976). A class of Bianchi type-VI cosmological models with electromagnetic field. *The Astrophysical Journal*, **204**, 322-329.
- Einstein, A. (1915). Zur allgemeinen Relativitätstheorie. *Preuss. Akad. Wiss. Berlin, Sitzber.*, **778**.
- Einstein, A. (1917). Kosmologische Betrachtungen zur allgemeinen Relativitätstheorie. *Preuss. Akad. Wiss. Berlin, Sitzber.*, **141**.
- Eisenstein, D. J., Zehavi, I., Hogg, D. W., Scoccamarro, R., Blanton, M. R., Nichol, R. C., ... & York, D. G. (2005). Detection of the baryon acoustic peak in the large-scale correlation function of SDSS luminous red galaxies. *The Astrophysical Journal*, **633**(2), 560.

- Ellis, G. F., & MacCallum, M. A. (1969). A class of homogeneous cosmological models. *Communications in Mathematical Physics*, **12(2)**, 108-141.
- Elmardi, M., Abebe, A., & Tekola, A. (2016). Chaplygin-gas solutions of  $f(R)$  gravity. *International Journal of Geometric Methods in Modern Physics*, **13(10)**, 1650120.
- Fabris, J. C., Goncalves, S. V. B., & Ribeiro, R. D. (2006). Bulk viscosity driving the acceleration of the Universe. *General Relativity and Gravitation*, **38(3)**, 495-506.
- Farajollahi, H., Salehi, A., & Nasiri, M. (2011). Stability analysis and observational constraints in scalar tensor theory. *Journal of Cosmology and Astroparticle Physics*, **2011(11)**, 018.
- Feng, B., Li, M., Piao, Y. S., & Zhang, X. (2006). Oscillating quintom and the recurrent universe. *Physics Letters B*, **634(2-3)**, 101-105.
- Friedmann, A. (1922). 125. on the curvature of space. *Zeitschrift für Physik*, **10**, 377-386.
- Frieman, J. A., Turner, M. S., & Huterer, D. (2008). Dark energy and the accelerating universe. *Annu. Rev. Astron. Astrophys.*, **46**, 385-432.
- Ftaclas, C., & Cohen, J. M. (1979). Tilted Bianchi type-III cosmologies. *Physical Review D*, **19(4)**, 1051.
- Ghate, H. R., & Sontakke, A. S. (2013). Bianchi type-IX dark energy model in Sen-Dunn scalar-tensor theory of gravitation. *Prespacetime Journal*, **5**, 619.
- Ghate, H. R., & Sontakke, A. S. (2014). Bianchi type-IX magnetized dark energy model in saez-ballester theory of gravitation. *International Journal of Astronomy and Astrophysics*, **4(1)**.



- Gorini, V., Kamenshchik, A., & Moschella, U. (2003). Can the Chaplygin gas be a plausible model for dark energy?. *Physical Review D*, **67(6)**, 063509.
- Guo, Z. K., Piao, Y. S., Zhang, X., & Zhang, Y. Z. (2005). Cosmological evolution of a quintom model of dark energy. *Physics Letters B*, **608(3-4)**, 177-182.
- Guo, Z. K., & Zhang, Y. Z. (2007). Cosmology with a variable Chaplygin gas. *Physics Letters B*, **645(4)**, 326-329.
- Halford, W. D. (1972). Scalar-tensor theory of gravitation in a Lyra manifold. *Journal of Mathematical Physics*, **13(11)**, 1699-1703.
- Harrison, E. R. (1973). Origin of magnetic fields in the early universe. *Physical Review Letters*, **30(5)**, 188.
- Hatkar, S. P., Wadale, C. D., & Katore, S. D. (2019). Bianchi Type-III Cosmological Model with Bulk Viscosity and Heat Flux in Modified  $f(R, T)$  Theory of Gravitation. *Bulg. J. Phys*, **46**, 107-116.
- Harko, T., Lobo, F. S., Nojiri, S. I., & Odintsov, S. D. (2011).  $f(R, T)$  gravity. *Physical Review D*, **84(2)**, 024020.
- Hegazy, E. A. (2019). Bianchi type VI cosmological model in self-creation theory. *Iranian Journal of Science and Technology, Transactions A: Science*, **43(2)**, 663-669.
- Hegazy, E. A., & Rahaman, F. (2020). Bianchi type- $VI_0$  cosmological model with electromagnetic field and variable deceleration parameter in general theory of relativity. *Indian Journal of Physics*, **94(11)**, 1847-1852.
- Hoftuft, J., Eriksen, H. K., Banday, A. J., Gorski, K. M., Hansen, F. K., & Lilje, P. B. (2009). Increasing evidence for hemispherical power asymmetry in the five-year WMAP data. *The Astrophysical Journal*, **699(2)**, 985.

- Hu, W., & Dodelson, S. (2002). Cosmic microwave background anisotropies. *Annual Review of Astronomy and Astrophysics*, **40**(1), 171-216.
- Hubble, E. (1929). A relation between distance and radial velocity among extragalactic nebulae. *Proceedings of the National Academy of Sciences(USA)*, **15**, 168.
- Jaffe, T. R., Banday, A. J., Eriksen, H. K., Górski, K. M., & Hansen, F. K. (2005). Evidence of vorticity and shear at large angular scales in the WMAP data: a violation of cosmological isotropy?. *The Astrophysical Journal*, **629**(1), L1.
- Johri, V. B., & Desikan, K. (1994). Cosmological models with constant deceleration parameter in Brans-Dicke theory. *General Relativity and Gravitation*, **26**(12), 1217-1232.
- Johri, V. B., & Sudharsan, R. (1989). BD-FRW cosmology with bulk viscosity. *Australian Journal of Physics*, **42**(2), 215-222.
- Jungman, G., Kamionkowski, M., & Griest, K. (1996). Supersymmetric dark matter. *Physics Reports*, **267**(5-6), 195-373.
- Kantowski, R., & Sachs, R. K. (1966). Some spatially homogeneous anisotropic relativistic cosmological models. *Journal of Mathematical Physics*, **7**(3), 443-446.
- King, D. H. (1991). Gravity-wave insights to Bianchi type-IX universes. *Physical Review D*, **44**(8), 2356.
- Kiran, M., & Reddy, D. R. K. (2013). Non-existence of Bianchi type-III bulk viscous string cosmological model in  $f(R, T)$  gravity. *Astrophysics and Space Science*, **346**(2), 521-524.
- Kiran, M., Reddy, D. R. K., & Rao, V. U. M. (2015). Bianchi type-III minimally interacting holographic dark energy model with linearly varying deceleration parameter in Brans-Dicke theory. *Astrophysics and Space Science*, **360**(2), 1-5.

- Knop, R. A., Aldering, G., Amanullah, R., Astier, P., Blanc, G., Burns, M. S., ... & Supernova Cosmology Project. (2003). New constraints on  $\Omega_M$ ,  $\Omega_\Lambda$ , and  $w$  from an independent set of 11 high-redshift supernovae observed with the Hubble Space Telescope. *The Astrophysical Journal*, **598(1)**, 102.
- Korunur, M. (2019). Tsallis holographic dark energy in Bianchi type-III spacetime with scalar fields. *Modern Physics Letters A*, **34(37)**, 1950310.
- Krori, K. D., & Nandy, D. (1977). Birkhoff's theorem and scalar-tensor theories of gravitation. *Journal of Physics A: Mathematical and General*, **10(6)**, 993.
- Kristian, J., & Sachs, R. K. (1966). Observations in cosmology. *The Astrophysical Journal*, **143**, 379.
- Latifi, A., Musette, M., & Conte, R. (1994). The Bianchi IX (mixmaster) cosmological model is not integrable. *Physics Letters A*, **194(1-2)**, 83-92.
- Leon, G., Paliathanasis, A., & Morales-Martínez, J. L. (2018). The past and future dynamics of quintom dark energy models. *The European Physical Journal C*, **78(9)**, 1-22.
- Li, M., Li, X. D., Wang, S., & Wang, Y. (2011). Dark energy. *Communications in theoretical physics*, **56(3)**, 525.
- Liddle, A. R., & Lyth, D. H. (1993). The Cold dark matter density perturbation. *Physics Reports*, **231(1-2)**, 1-105.
- Lorenz, D. (1982). Tilted electromagnetic Bianchi type-III cosmological model. *Astrophysics and Space Science*, **85(1)**, 63-67.
- Lorenz, D. (1982). Exact Bianchi type- $VI_0$  cosmological models with matter and an electromagnetic field. *Astrophysics and Space Science*, **85(1)**, 69-74.

- Ludwick, K. J. (2017). The viability of phantom dark energy: A review. *Modern Physics Letters A*, **32(28)**, 1730025.
- Lyra, G. (1951). Über eine modifikation der riemannschen geometrie. *Mathematische Zeitschrift*, **54(1)**, 52-64.
- Mahanta, K. L., Biswal, A. K., & Sahoo, P. K. (2014). Bianchi type-III dark energy models with constant deceleration parameter in self-creation cosmology. *Canadian Journal of Physics*, **92(4)**, 295-301.
- Malquarti, M., Copeland, E. J., & Liddle, A. R. (2003). K-essence and the coincidence problem. *Physical Review D*, **68(2)**, 023512.
- Matravers, D. R., & Tsagas, C. G. (2000). Cosmic magnetism, curvature, and the expansion dynamics. *Physical Review D*, **62(10)**, 103519.
- Melvin, M. A. (1975). Homogeneous axial cosmologies with electromagnetic fields and dust. *Ann. NY Acad. Sci.:(United States)*,**262**.
- Mohanty, G., & Mishra, B. (2002). Bianchi type- $VI_1$  cosmological model in scale-invariant theory. *Czechoslovak journal of physics*, **52(6)**, 765-773.
- Mishra, R. K., & Chand, A. (2016a). Cosmological models in alternative theory of gravity with bilinear deceleration parameter. *Astrophysics and Space Science*, **361(8)**, 1-10.
- Mishra, R. K., Chand, A., & Pradhan, A. (2016b). Dark energy models in  $f(R, T)$  theory with variable deceleration parameter. *International Journal of Theoretical Physics*, **55(2)**, 1241-1256.
- Mishra, R. K., & Chand, A. (2017). A comparative study of cosmological models in alternative theory of gravity with LVDP & BVDP. *Astrophysics and Space Science*, **362(8)**, 1-11.

- Mishra, R. K., & Chand, A. (2020). Cosmological models in Sáez-Ballester theory with bilinear varying deceleration parameter. *Astrophysics and Space Science*, **365(4)**, 1-9.
- Mishra, R. K., Dua, H., & Chand, A. (2018). Bianchi-III cosmological model with BVDP in modified  $f(R, T)$  theory. *Astrophysics and Space Science*, **363(6)**, 1-8.
- Mishra, R. K., & Dua, H. (2019). Bulk viscous string cosmological models in Saez-Ballester theory of gravity. *Astrophysics and Space Science*, **364(11)**, 1-12.
- Mishra, B., & Sahoo, P. K. (2014). Bianchi type- $VI_1$  cosmological model with wet dark fluid in scale invariant theory of gravitation. *Astrophysics and Space Science*, **349(1)**, 491-499.
- Misner, C. W. (1969). Mixmaster universe. *Physical Review Letters*, **22(20)**, 1071.
- Mohanty, G., & Mishra, B. (2002). Bianchi type- $VI_1$  cosmological model in scale-invariant theory. *Czechoslovak journal of physics*, **52(6)**, 765-773.
- Mukherjee, B. (2003). Cylindrically symmetric solution of relativistic magneto-fluid in Sen-Dunn theory of gravitation. *Indian Journal of Pure and Applied Mathematics*, **34(7)**, 1121-1127.
- Mukherjee, B. (2004). Static spherically symmetric anisotropic fluid distribution in presence of electromagnetic field in the Sen-Dunn theory of gravitation. *Acta Physica Hungarica Series A, Heavy Ion Physics*, **21(1)**, 39-47.
- Naidu, R. L., Aditya, Y., & Reddy, D. R. K. (2019). Bianchi type-V dark energy cosmological model in general relativity in the presence of massive scalar field. *Heliyon*, **5(5)**, e01645.
- Naidu, R. L., Satyanarayana, B., & Reddy, D. R. K. (2012). Bianchi type-III dark energy model in a Saez-Ballester scalar-tensor theory. *International Journal of Theoretical Physics*, **51(9)**, 2857-2862.

- Narlikar, J. V. (2002). *An introduction to cosmology*. Cambridge University Press.
- Navarro, J. F. (1996). The structure of cold dark matter halos. *In Symposium-international astronomical union, Cambridge University Press.*, **171**, 255-258.
- Nojiri, S. I., & Odintsov, S. D. (2006). Unifying phantom inflation with late-time acceleration: Scalar phantom-non-phantom transition model and generalized holographic dark energy. *General Relativity and Gravitation*, **38(8)**, 1285-1304.
- Padmanabhan, T. (2002). Accelerated expansion of the universe driven by tachyonic matter. *Physical Review D*, **66(2)**, 021301.
- Padmanabhan, T. (2003). Cosmological constant—the weight of the vacuum. *Physics Reports*, **380(5-6)**, 235-320.
- Page, L., Barnes, C., Hinshaw, G., Spergel, D. N., Weiland, J. L., Wollack, E., ... & Wright, E. L. (2003). First-year wilkinson microwave anisotropy probe (WMAP) observations: Beam profiles and window functions. *The Astrophysical Journal Supplement Series*, **148(1)**, 39.
- Panotopoulos, G., & Rincón, Á. (2018). Growth index and statefinder diagnostic of oscillating dark energy. *Physical Review D*, **97(10)**, 103509.
- Patel, L. K., & Koppal, S. S. (1991). Some Bianchi type- $VI_0$  viscous fluid cosmological models. *The ANZIAM Journal*, **33(1)**, 77-84.
- Pawar, D. D., & Shahare, S. P. (2019). Dynamics of tilted Bianchi type-III cosmological model in  $f(R, T)$  gravity. *Journal of Astrophysics and Astronomy*, **40(4)**, 1-7.
- Peebles, P. J. E., & Ratra, B. (1988). Cosmology with a time-variable cosmological 'constant'. *The Astrophysical Journal*, **325**, L17-L20.

- Peebles, P. J. E., & Ratra, B. (2003). The cosmological constant and dark energy. *Reviews of modern physics*, **75(2)**, 559.
- Perlmutter, S., Aldering, G., Valle, M. D., Deustua, S., Ellis, R. S., Fabbro, S., ... & Walton, N. (1998). Discovery of a supernova explosion at half the age of the Universe. *Nature*, **391(6662)**, 51-54.
- Perlmutter, S., Aldering, G., Goldhaber, G., Knop, R. A., Nugent, P., Castro, P. G., ...& Supernova Cosmology Project. (1999). Measurements of  $\Omega$  and  $\Lambda$  from 42 high-redshift supernovae. *The Astrophysical Journal*, **517(2)**, 565.
- Piazza, F., & Tsujikawa, S. (2004). Dilatonic ghost condensate as dark energy. *Journal of Cosmology and Astroparticle Physics*, **2004(07)**, 004.
- Pospelov, M., Ritz, A., & Voloshin, M. (2008). Secluded WIMP dark matter. *Physics Letters B*, **662(1)**, 53-61.
- Pradhan, A., Srivastav, S. K., & Yadav, M. K. (2005). Some homogeneous Bianchi type-IX viscous fluid cosmological models with a varying  $\Lambda$ . *Astrophysics and Space Science*, **298(3)**, 419-432.
- Pradhan, A., Lata, S., & Amirhashchi, H. (2010). Massive string cosmology in Bianchi type-III space-time with electromagnetic field. *Communications in Theoretical Physics*, **54(5)**, 950.
- Pradhan, A., & Amirhashchi, H. (2011). Dark energy model in anisotropic Bianchi type-III space-time with variable EoS parameter. *Astrophysics and Space Science*, **332(2)**, 441-448.
- Pradhan, A., Jaiswal, R., Jotania, K., & Khare, R. K. (2012). Dark energy models with anisotropic fluid in Bianchi type- $VI_0$  space-time with time dependent deceleration parameter. *Astrophysics and Space Science*, **337(1)**, 401-413.

- Pudritz, R. E., & Silk, J. (1989). The origin of magnetic fields and primordial stars in protogalaxies. *The Astrophysical Journal*, **342**, 650-659.
- Raju, K. D., Aditya, Y., Rao, V. U. M., & Reddy, D. R. K. (2020). Bianchi type-III dark energy cosmological model with massive scalar meson field. *Astrophysics and Space Science*, **365(2)**, 1-7.
- Rao, V. U. M., & Neelima, D. (2013). Bianchi type- $VI_0$  perfect fluid cosmological model in a modified theory of gravity. *Astrophysics and Space Science*, **345(2)**, 427-430.
- Raychaudhuri, A. K. (1979). Theoretical cosmology. *Oxford Studies in Physics*.
- Reddy, D. R. K. (1973). On Birkhoff's theorem in scalar-tensor theory of gravitation. *Journal of Physics A: Mathematical, Nuclear and General*, **6(12)**, 1867.
- Reddy, D. R. K. (1977). Static plane-symmetric solutions in Brans-Dicke and Sen-Dunn theories of gravitation. *Journal of Physics A: Mathematical and General*, **10(1)**, 55.
- Reddy, D. R. K., Anitha, S., & Umadevi, S. (2016). Anisotropic holographic dark energy model in Bianchi type- $VI_0$  universe in a scalar-tensor theory of gravitation. *Astrophysics and Space Science*, **361(10)**, 1-7.
- Reddy, D. R. K., & Naidu, R. L. (2007). Bianchi type-IX cosmic strings in a scalar-tensor theory of gravitation. *Astrophysics and Space Science*, **312(1)**, 99-102.
- Reddy, D. R., Santikumar, R., & Naidu, R. L. (2012). Bianchi type-III cosmological model in  $f(R, T)$  theory of gravity. *Astrophysics and Space Science*, **342(1)**, 249-252.
- Reddy, D. R. K., Santhi Kumar, R., & Pradeep Kumar, T. V. (2013). Bianchi type-III dark energy model in  $f(R, T)$  gravity. *International Journal of Theoretical Physics*, **52(1)**, 239-245.



- Reddy, D. R. K., & Venkateswarlu, R. (1987). Birkhoff-type theorem in the scale-covariant theory of gravitation. *Astrophysics and space science*, **136(1)**, 191-194.
- Ribeiro, M. B., & Sanyal, A. K. (1987). Bianchi  $VI_0$  viscous fluid cosmology with magnetic field. *J. Math. Phys.(NY);(United States)*, **28(3)**.
- Riess, A. G., Filippenko, A. V., Challis, P., Clocchiatti, A., Diercks, A., Garnavich, P. M., ... & Tonry, J. (1998). Observational evidence from supernovae for an accelerating universe and a cosmological constant. *The Astronomical Journal*, **116(3)**, 1009.
- Riess, A. G., Nugent, P. E., Gilliland, R. L., Schmidt, B. P., Tonry, J., Dickinson, M., ...& Veilleux, S. (2001). The farthest known supernova: support for an accelerating universe and a glimpse of the epoch of deceleration. *The Astrophysical Journal*, **560(1)**, 49.
- Riess, A. G., Strolger, L. G., Tonry, J., Casertano, S., Ferguson, H. C., Mobasher, B., ... & Tsvetanov, Z. (2004). Type Ia supernova discoveries at  $z > 1$  from the Hubble Space Telescope: Evidence for past deceleration and constraints on dark energy evolution. *The Astrophysical Journal*, **607(2)**, 665.
- Romano, V., & Pavón, D. (1994). Dissipative effects in Bianchi type-III cosmologies. *Physical Review D*, **50(4)**, 2572.
- Roy, A. R., & Chatterjee, B. (1980). Cylindrically symmetric empty space-time solutions of SEN-DUNN theory. *Acta Physica Academiae Scientiarum Hungaricae*, **48(4)**, 383-389.
- Roy, A. R., & Chatterjee, B. (1981). Plane-symmetric static charged-dust distribution in Sen-Dunn theory. *Indian Journal of Pure & Applied Mathematics*, **12(5)**, 659-663.

- Roy, S. R., Narain, S., & Singh, J. P. (1985a). Bianchi type I cosmological models with perfect fluid and magnetic field. *Australian Journal of Physics*, **38(2)**, 239-248.
- Roy, S. R., Singh, J. P., & Narain, S. (1985b). Bianchi type- $VI_0$  perfect fluid cosmological models with magnetic field. *Astrophysics and space science*, **111(2)**, 389-397.
- Sahni, V., Saini, T. D., Starobinsky, A. A., & Alam, U. (2003). Statefinder-a new geometrical diagnostic of dark energy. *Journal of Experimental and Theoretical Physics Letters*, **77(5)**, 201-206.
- Sahni, V., & Starobinsky, A. (2000). The case for a positive cosmological  $\Lambda$ -term. *International Journal of Modern Physics D*, **9(04)**, 373-443.
- Sahni, V., & Starobinsky, A. (2006). Reconstructing dark energy. *International Journal of Modern Physics D*, **15(12)**, 2105-2132.
- Sahoo, P. K., & Mishra, B. (2015). Higher-dimensional Bianchi type-III universe with strange quark matter attached to string cloud in general relativity. *Turkish Journal of Physics*, **39(1)**, 43-53.
- Sahoo, P. K., Nath, A., & Sahu, S. K. (2017). Bianchi type-III string cosmological model with bulk viscous fluid in Lyra geometry. *Iranian Journal of Science and Technology, Transactions A: Science*, **41(1)**, 243-248.
- Sahoo, P. K., Sahoo, P., Bishi, B. K., & Aygün, S. (2018). Magnetized strange quark matter in  $f(R, T)$  gravity with bilinear and special form of time varying deceleration parameter. *New Astronomy*, **60**, 80-87.
- Saleem, R., & Imtiaz, M. J. (2020). Dynamical study of interacting Ricci dark energy model using Chevallier-Polarsky-Lindertype parametrization. *Classical and Quantum Gravity*, **37(6)**, 065018.

Samanta, G. C. (2013). Bianchi type-III cosmological models with anisotropic dark energy (DE) in lyra geometry. *International Journal of Theoretical Physics*, **52(10)**, 3442-3456.

Santhi, M. V., Rao, V. U. M., & Aditya, Y. (2017). Bianchi type-VI<sub>0</sub> modified holographic Ricci dark energy model in a scalar–tensor theory. *Canadian Journal of Physics*, **95(2)**, 179-183.

Santhi, M. V., Rao, V. U. M., & Aditya, Y. (2017). Anisotropic generalized ghost pilgrim dark energy model in general relativity. *International Journal of Theoretical Physics*, **56(2)**, 362-371.

Santhi, M. V., Rao, V. U. M., Gusu, D. M., & Aditya, Y. (2018). Bianchi type-III holographic dark energy model with quintessence. *International Journal of Geometric Methods in Modern Physics*, **15(09)**, 1850161.

Santhi, M. V., & Sobhanbabu, Y. (2020). Bianchi type-III Tsallis holographic dark energy model in Seaz-Ballester theory of gravitation. *The European Physical Journal C*, **80(12)**, 1198.

Sarkar, S. (2014). Holographic dark energy model with linearly varying deceleration parameter and generalised Chaplygin gas dark energy model in Bianchi type-I universe. *Astrophysics and Space Science*, **349(2)**, 985-993.

Satish, J., & Venkateswarlu, R. (2019). Anisotropic Bianchi Type-VI<sub>0</sub> Two Fluid Cosmological Model Coupled with Massless Scalar Field and Time-Varying G and  $\Lambda$ . *Bulgarian Journal of Physics*, **46(1)**.

Schmidt, B. P., Suntzeff, N. B., Phillips, M. M., Schommer, R. A., Clocchiatti, A., Kirshner, R. P., ... & Ciardullo, R. (1998). The high-Z supernova search: measuring cosmic deceleration and global curvature of the universe using type Ia supernovae. *The Astrophysical Journal*, **507(1)**, 46.

- Sen, D. K. (1957). A static cosmological model. *Zeitschrift für Physik*, **149(3)**, 311-323.
- Sen, A. (2002). Tachyon matter. *Journal of High Energy Physics*, **2002(07)**, 065.
- Sen, D. K., & Dunn, K. A. (1971). A scalar-tensor theory of gravitation in a modified Riemannian manifold. *Journal of Mathematical Physics*, **12(4)**, 578-586.
- Shamir, M. F. (2011). Plane symmetric vacuum Bianchi type-III cosmology in  $f(R)$  gravity. *International Journal of Theoretical Physics*, **50(3)**, 637-643.
- Sharif, M., & Zubair, M. (2010). Effects of electromagnetic field on the dynamics of bianchi type- $VI_0$  universe with anisotropic dark energy. *International Journal of Modern Physics D*, **19(12)**, 1957-1972.
- Sharif, M., & Kausar, H. R. (2011). Anisotropic fluid and Bianchi type-III model in  $f(R)$  gravity. *Physics Letters B*, **697(1)**, 1-6.
- Sharif, M., & Kausar, H. R. (2011). Non-vacuum solutions of Bianchi type- $VI_0$  universe in  $f(R)$  gravity. *Astrophysics and Space Science*, **332(2)**, 463-471.
- Sharma, U. K., Zia, R., Pradhan, A., & Beesham, A. (2019). Stability of LRS Bianchi type-I cosmological models in  $f(R, T)$ -gravity. *Research in Astronomy and Astrophysics*, **19(4)**, 055.
- Sharma, S., & Poonia, L. (2021). Cosmic inflation in Bianchi Type IX space with bulk viscosity. *Advances in Mathematics: Scientific Journal*, **10(1)**, 527-534.
- Singh, C. P. (2008). Bulk viscous cosmology in early universe. *Pramana*, **71(1)**, 33-48.
- Singh, J. P. (2008). A cosmological model with both deceleration and acceleration. *Astrophysics Space Sciences*, **318**, 103-107

- Singh, C. P., & Kumar, P. (2016). Statefinder diagnosis for holographic dark energy models in modified  $f(R, T)$  gravity. *Astrophysics and Space Science*, **361(5)**, 1-9.
- Singh, T., & Rai, L. N. (1983). Scalar-tensor theories of gravitation: foundations and prospects. *General Relativity and Gravitation*, **15(9)**, 875-902.
- Singh, J. K., & Ram, S. (1996). String cosmological models of Bianchi type-III. *Astrophysics and Space Science*, **246(1)**, 65-72.
- Singha, A. K., & Debnath U. (2009). Accelerating Universe with a special form of decelerating paramter. *International Journal of Theoretical Physics*, **48**, 351-356
- Smith, P. F., & Lewin, J. D. (1990). Dark matter detection. *Physics Reports*, **187(5)**, 203-280.
- Sofuoğlu, D. (2016). Rotating and expanding Bianchi type-IX model in  $f(R, T)$  theory of gravity. *Astrophysics and Space Science*, **361(1)**, 1-7.
- Spergel, D. N., & Steinhardt, P. J. (2000). Observational evidence for self-interacting cold dark matter. *Physical review letters*, **84(17)**, 3760.
- Steigman, G. (2007). Primordial nucleosynthesis in the precision cosmology era. *Annual Review of Nuclear and Particle Science*, **57**, 463-491
- Steinhardt, P. J., Wang, L., & Zlatev, I. (1999). Cosmological tracking solutions. *Physical Review D*, **59(12)**, 123504.
- Taub, A. H. (1951). Empty space-times admitting a three parameter group of motions. *Annals of Mathematics*, 472-490.
- Tegmark, M., Strauss, M. A., Blanton, M. R., Abazajian, K., Dodelson, S., Sandvik, H., ... & York, D. G. (2004). Cosmological parameters from SDSS and WMAP. *Physical review D*, **69(10)**, 103501.

- Thorne, K. S. (1967). Primordial element formation, primordial magnetic fields, and the isotropy of the universe. *The Astrophysical Journal*, **148**, 51.
- Tian, S. X., & Zhu, Z. H. (2021). Early dark energy in k-essence. *Physical Review D*, **103(4)**, 043518.
- Tikekar, R., & Patel, L. K. (1992). Some exact solutions of string cosmology in Bianchi III space-time. *General Relativity and Gravitation*, **24(4)**, 397-404.
- Tiwari, R. K., Agrawal, A. K., & Shukla, B. K. (2016). Bianchi type- $VI_0$  cosmological model with variable deceleration parameter. *Prespacetime Journal*, **7(5)**.
- Tiwari, R. K., Beesham, A., & Shukla, B. K. (2016). Behaviour of the cosmological model with variable deceleration parameter. *The European Physical Journal Plus*, **131(12)**, 1-9.
- Tiwari, R. K., & Beesham, A. (2018). Anisotropic model with decaying cosmological term. *Astrophysics Space Science*, **363**, 234.
- Tripathy, S. K., Sahu, S. K., & Routray, T. R. (2008). String cloud cosmologies for Bianchi type-III models with electromagnetic field. *Astrophysics and Space Science*, **315(1)**, 105-110.
- Tsagas, C. G., & Barrow, J. D. (1997). A gauge-invariant analysis of magnetic fields in general-relativistic cosmology. *Classical and Quantum Gravity*, **14(9)**, 2539.
- Tsujikawa, S. (2013). Quintessence: a review. *Classical and Quantum Gravity*, **30(21)**, 214003.
- Tyagi, A., & Chhajed, D. (2012). Homogeneous anisotropic Bianchi type-IX cosmological model for perfect fluid distribution with electromagnetic field. *American journal of Mathematics and statistics*, **2(3)**, 19-21.

- Uggla, C., & Zur-Muhlen, H. (1990). Compactified and reduced dynamics for locally rotationally symmetric Bianchi type IX perfect fluid models. *Classical and Quantum Gravity*, **7(8)**, 1365.
- Umadevi, S., & Ramesh, G. (2015). Minimally interacting holographic dark energy model in Bianchi type-III universe in Brans-Dicke theory. *Astrophysics and Space Science*, **359(2)**, 1-5.
- Upadhaya, R. D., & Dave, S. (2008). Some magnetized bianchi type-III massive string cosmological models in general relativity. *Brazilian Journal of Physics*, **38**, 615-620.
- Varshney, G., Sharma, U. K., & Pradhan, A. (2019). Statefinder diagnosis for interacting Tsallis holographic dark energy models with  $\omega - \omega'$  pair. *New Astronomy*, **70**, 36-42.
- Vidya Sagar, T., Purnachandra Rao, C., Bhuvana Vijaya, R., & Reddy, D. R. K. (2014). Bianchi type-III bulk viscous string cosmological model in Brans-Dicke theory of gravitation. *Astrophysics and Space Science*, **349(1)**, 479-483.
- Visser, M., & Barcelo, C. (2000). Energy conditions and their cosmological implications. *In Cosmo-99* (pp. 98-112).
- Venkateswarlu, R., Satish, J., & Kumar, K. P. (2011). Some Bianchi type-I Cosmic Strings in a Scalar-Tensor Theory of Gravitation. *EJTP*, **8(25)**, 353-360.
- Venkateswarlu, R., Satish, J., & Kumar, K. P. (2013a). Bianchi Type- $VI_0$  String Cosmological Models in a New Scalar-Tensor Theory of Gravitation. *Prespacetime Journal*, **4(4)**.
- Venkateswarlu, R., Satish, J., & Pavan Kumar, K. (2013b). Higher Dimensional FRW String Cosmological Models in a New Scalar-tensor Theory of Gravitation. *The African Review of Physics*, **8**.

- Venkateswarlu, R., Satish, J., & Kumar, K. P. (2013c). LRS Bianchi Type-III Massive String Cosmological Models in Scalar Theory of Gravitation. *Prespacetime Journal*, **4(3)**.
- Waller, S. M. (1984). Bianchi type-IX electromagnetic universes. *Physical Review D*, **29(2)**, 176.
- Weinberg, S. (1989). The cosmological constant problem. *Reviews of modern physics*, **61(1)**, 1.
- Weinberg, S. (2015). *To explain the world: The discovery of modern science*. Penguin UK
- Wei, H., & Cai, R. G. (2005). Cosmological evolution of “hessence” dark energy and avoidance of the big rip. *Physical Review D*, **72(12)**, 123507.
- Wei, H., Cai, R. G., & Zeng, D. F. (2005). Hessian: a new view of quintom dark energy. *Classical and Quantum Gravity*, **22(16)**, 3189.
- Weyl, H. (1918). Reine infinitesimalgeometrie. *Mathematische Zeitschrift*, **2(3)**, 384-411.
- Xing-Xiang, W. (2005). Bianchi type-III string cosmological model with bulk viscosity in general relativity. *Chinese Physics Letters*, **22(1)**, 29.
- Xing-Xiang, W. (2006). Bianchi type-III string cosmological model with bulk viscosity and magnetic field. *chinese physics letters*, **23(7)**, 1702.
- Yadav, A. K. (2011). Some anisotropic dark energy models in Bianchi type-V space-time. *Astrophysics and Space Science*, **335(2)**, 565-575.
- Yadav, M. K., Rai, A., & Pradhan, A. (2007). Some Bianchi Type-III string cosmological models with bulk viscosity. *International Journal of Theoretical Physics*, **46(11)**, 2677-2687.



Yadav, A. K., & Yadav, L. (2011). Bianchi type-III anisotropic dark energy models with constant deceleration parameter. *International Journal of Theoretical Physics*, **50(1)**, 218-227.

Zeldovich, Y. B., Ruzmaikin, A. A., & Sokolov, D. D. (1983). Magnetic fields in astrophysics. *New York*, 3.

Zia, R., Maurya, D. C., & Pradhan, A. (2018). Transit dark energy string cosmological models with perfect fluid in F (R, T)-gravity. *International Journal of Geometric Methods in Modern Physics*, **15(10)**, 1850168.

Zimdahl, W., Pavon, D., & Chimento, L. P. (2001). Interacting quintessence. *Physics Letters B*, **521(3-4)**, 133-138.

## Annexure-I

### **The List of Published papers are:**

1. Basumatay, D., & Dewri, M. (2021). Bianchi Type- $VI_0$  Cosmological Model with Special Form of Scale Factor in Sen-Dunn Theory of Gravitation. *Journal of Scientific Research*, 13(1), 137-143.
2. Basumatay, D., & Dewri, M. (2022). Magnetized Bianchi model with time-dependent deceleration parameter. *Indian Journal of Science and Technology*, 15(35), 1703-1711.

## Annexure-II

### **The List of Seminar papers presented are:**

1. Bianchi Type-IX Cosmological Model in Sen-Dunn Theory of Gravitation, *National Conference on Emerging Trends in Advanced Mathematical Sciences and its Inter-Disciplinary Areas*, Organized by Department of Mathematical Sciences, Bodoland University, Kokrajhar, India on 21-22 February 2020.
2. Bianchi Type- $VI_0$  Cosmological model with special form of scale factor in Sen-Dunn theory of Gravitation, *National conference on Advanced Mathematical & Applied Sciences*, Organized by Department of Mathematical Sciences, Bodoland University, Kokrajhar, India, on 1-2 July 2020
3. Magnetized Bianchi type- $VI_0$  dark energy model with electromagnetic field, *National Conference on Science & Technology for Sustainable Development*, Organized by Science College, Kokrajhar, Assam, India, on 9-10th September 2022.

## Bianchi Type- $VI_0$ Cosmological Model with Special Form of Scale Factor in Sen-Dunn Theory of Gravitation

D. Basumatary\*, M. Dewri

Department of Mathematical Sciences, Bodoland University, Kokrajhar, 783370, India

Received 2 August 2020, accepted in final revised form 13 October 2020

### Abstract

A Bianchi Type- $VI_0$  cosmological model with a special form of scale factor is studied. Einstein field equations in Sen-Dunn theory are obtained and solved for exact solutions. This solution gives a scenario of the dark energy model which tends to a  $\Lambda$ CDM model. The physical and geometrical properties are also obtained and analyzed with the present day observations.

*Keywords:* Sen-Dunn theory; Dark energy;  $\Lambda$ CDM model; Bianchi type- $VI_0$ .

© 2021 JSR Publications. ISSN: 2070-0237 (Print); 2070-0245 (Online). All rights reserved.  
doi: <http://dx.doi.org/10.3329/jsr.v13i1.48479> J. Sci. Res. **13** (1), 137-143 (2021)

### 1. Introduction

Einstein General Theory of Relativity is one of the most essential and dynamic phenomena, benefiting in describing and creating models to understand the universe better. Concerning Albert Einstein's theory of gravitation, many other authors developed an alternative theory to Einstein's theory of gravitation developed in recent decades. Like an alternative theory of gravitation by Brans-Dickie [1], the tensor field is identified by the metric tensor of Riemannian geometry and the scalar field, which is alien to the geometry. Sen and Dunn [2] proposed a theory of gravitation based on Lyra's geometry, where the field equation contains metric tensor  $g_{ij}$  and scalar function  $\varphi$  which is considered as special character of the theory are both geometrized. Dunn [4] proposed the scalar-tensor theory of gravitation using a non-Riemannian geometry in which both the metric tensor and the scalar function have an unambiguous geometric interpretation with a function  $\varphi = \varphi(x^i)$  where  $x^i$  is the coordinate in a four dimensional Riemannian manifold. Birkhoff's theorem and spherically symmetric static conformally flat solutions are studied using the Sen-Dunn scalar-tensor theory by Reddy [3,5]. Roy and Chatterjee [6] investigated symmetrically charged dust distribution in the Sen-Dunn theory of gravitation. Also, many authors like Singh and Rai [7], Mukherjee [8], Reddy and Naidu [9], Venkateswarlu *et al.* [10,11] had investigated various aspects of cosmological models

---

\*Corresponding author: [flindstif@gmail.com](mailto:flindstif@gmail.com)

in a different context with the Sen-Dunn scalar-tensor theory of gravitation. Ghate and Sontakke [12] studied Bianchi type-IX dark energy model in the Sen-Dunn theory of gravitation. Patra [13] also studied the string cosmological model in a new scalar-tensor theory of gravitation. Bianchi type models give us the simplicity of the field equations and better understand the universe's anisotropy, which is relatively better than FRW isotropic models. Bianchi type models are homogeneous and anisotropic, giving a better image of the early stages of the present-day observation and the behavior in the different stages. Roy and Singh [14] investigated the Bianchi type  $VI_0$  cosmological model with a magnetic type with the free gravitational field. Patel and Koopar [15] studied viscous fluid with Bianchi type  $VI_0$  cosmological model. Bali *et al.* [16,17], Bali and Pradhan [18] studied the Bianchi type  $VI_0$  cosmological model relative to string theory. Many authors have mainly studied the aspects of dark energy cosmological models relative to the universe [19,20]. Mishra and Sahoo [21] investigated Bianchi type  $VI_1$  with the presence of wet, dark energy fluid cosmological model in scale-invariant theory. Anisotropic magnetized holographic Ricci dark energy cosmological models were discussed by Santhi *et al.* [22]. Also, gravity field equations are derived by Santhi *et al.* [23] with the help of a spatially homogeneous and anisotropic Bianchi type- $VI_h$  space-time in a bulk-viscous fluid, containing one-dimensional cosmic strings. Hegazy [24], Satish and Venkateswarlu [25] studied various aspects with Bianchi type  $VI_0$  cosmological models. Investigation of the Kantowski-Sachs cosmological model with bulk viscous and cosmic string in  $f(T)$  gravity framework is done by Bhojar *et al.* [26]. Recently, Dewri [27] studied the spatially homogeneous Robertson-Walker cosmological models with magnetized isotropic dark energy like fluid in the scalar-tensor theory of gravitation. Brahma and Dewri [28] also worked on a paper Bianchi Type – V Modified  $f(\mathbf{R}, \mathbf{T})$  model in Lyra manifold, that approach the  $\Lambda$ CDM (Lambda cold dark matter) model. Works of different researchers on Sen-Dunn and other scalar-tensor theory mentioned above are the motivation behind this work on accelerating and  $\Lambda$ CDM model. This paper has investigated the Bianchi type  $VI_0$  cosmological model having a variable EoS parameter and deceleration parameter in Sen-Dunn theory.

## 2. Metric and Field Equation

Bianchi type- $VI_0$  space-time metric is given by

$$ds^2 = -dt^2 + A^2 dx^2 + B^2 e^{2x} dy^2 + C^2 e^{-2x} dz^2 \quad (1)$$

where  $A, B, C$  are function of time  $t$ .

The generalization of EoS parameter of perfect fluid separately on each spatial axis by preserving the diagonal form and examining the diagonal form of the energy momentum tensor is.

$$T_{ij} = \text{diag}[T_{00}, T_{11}, T_{22}, T_{33}] \quad (2)$$

Thus, on parameterize it formulates

$$\begin{aligned} T_{ij} &= \text{diag}[\rho, -p_x, -p_y, -p_z] = \text{diag}[1, -w_x, -w_y, w_z]\rho \\ &= \text{diag}[1, -w, -(w + \delta), -(w + \gamma)]\rho \end{aligned} \quad (3)$$

where  $\rho$  is the energy density,  $p_x, p_y, p_z$  is the directional pressure,  $w_x, w_y, w_z$  are the directional EoS parameters along the  $x, y$  and  $z$  axis respectively,  $w$  is the considered as deviation free EoS parameter,  $\delta$  and  $\gamma$  are the skewness parameters. The parameters  $w, \delta$  and  $\gamma$  are essentially need not be constant and may be a function of time.

The field equation given by Sen-Dunn [1] for the combined scalar and tensor fields (in natural units  $c = 1, G = 1$ ) is

$$R_{ij} - \frac{1}{2}g_{ij}R = \omega\varphi^{-2}(\varphi_{,i}\varphi_{,j} - \frac{1}{2}g_{ij}\varphi_{,k}\varphi^{,k}) - \varphi^{-2}T_{ij} \tag{4}$$

where,  $\omega = \frac{3}{2}$ ,  $R_{ij}$ ,  $R$ ,  $g_{ij}$  and  $T_{ij}$  are the Ricci tensor, Ricci scalar, metric tensor and energy momentum tensor respectively.

The field equation (4) with (3) for the line element (1) gives rise to

$$\frac{\ddot{B}}{B} + \frac{\ddot{C}}{C} + \frac{\dot{B}\dot{C}}{BC} + \frac{1}{A^2} = \frac{-w\rho}{\varphi^2} + \frac{\omega}{2}\left(\frac{\dot{\varphi}}{\varphi}\right)^2 \tag{5}$$

$$\frac{\ddot{C}}{C} + \frac{\ddot{A}}{A} + \frac{\dot{A}\dot{C}}{AC} - \frac{1}{A^2} = \frac{-(w+\delta)\rho}{\varphi^2} + \frac{\omega}{2}\left(\frac{\dot{\varphi}}{\varphi}\right)^2 \tag{6}$$

$$\frac{\ddot{A}}{A} + \frac{\ddot{B}}{B} + \frac{\dot{A}\dot{B}}{AB} - \frac{1}{A^2} = \frac{-(w+\gamma)\rho}{\varphi^2} + \frac{\omega}{2}\left(\frac{\dot{\varphi}}{\varphi}\right)^2 \tag{7}$$

$$\frac{\dot{A}\dot{B}}{AB} + \frac{\dot{B}\dot{C}}{BC} + \frac{\dot{A}\dot{C}}{AC} - \frac{1}{A^2} = \frac{-\rho}{\varphi^2} + \frac{\omega}{2}\left(\frac{\dot{\varphi}}{\varphi}\right)^2 \tag{8}$$

$$\frac{\dot{C}}{C} - \frac{\dot{B}}{B} = 0 \tag{9}$$

The average scale factor for a Bianchi type  $VI_0$  space time is given as

$$a = (ABC)^{\frac{1}{3}} \tag{10}$$

Also, Hubble's parameter ( $H$ ), expansion scalar ( $\theta$ ), shear scalar ( $\sigma$ ), anisotropic parameter ( $A_m$ ), and deceleration parameter ( $q$ ) are defined as

$$H = \frac{\dot{a}}{a} \tag{11}$$

$$\theta = 3H \tag{12}$$

$$\sigma^2 = \frac{1}{2}\left(\sum_{i=1}^3 H_i^2 - \frac{1}{3}\theta^2\right) \tag{13}$$

$$A_m = \frac{1}{2}\sum_{i=1}^3\left(\frac{\Delta H_i}{H}\right)^2 = \frac{2\sigma^2}{3H^2} \tag{14}$$

$$q = \frac{-a\ddot{a}}{\dot{a}^2} = \frac{d}{dt}\left(\frac{1}{H}\right) - 1 \tag{15}$$

where,  $\Delta H_i = H_i - H$  ( $i = x, y, z$ ) represents the directional Hubble's parameters.

### 3. Solution of the Field Equation

From (9) we get by integrating

$$C = mB \tag{16}$$

where,  $m$  is a constant.

From (16), (7) and (6), the equality in the skewness parameters along the  $y$  and  $z$  axis is found.

So, the equations (5)-(9) reduces to

$$2\frac{\ddot{B}}{B} + \frac{\dot{B}^2}{B^2} + \frac{1}{A^2} = -\frac{w\rho}{\varphi^2} + \frac{\omega}{2}\left(\frac{\dot{\varphi}}{\varphi}\right)^2 \tag{17}$$

$$\frac{\ddot{A}}{A} + \frac{\ddot{B}}{B} + \frac{\dot{A}\dot{B}}{AB} - \frac{1}{A^2} = \frac{-(w+\delta)\rho}{\varphi^2} + \frac{\omega}{2}\left(\frac{\dot{\varphi}}{\varphi}\right)^2 \tag{18}$$

$$2\frac{\dot{A}\dot{B}}{AB} + \frac{\dot{B}}{B^2} - \frac{1}{A^2} = \frac{\rho}{\phi^2} - \frac{\omega}{2}\left(\frac{\dot{\phi}}{\phi}\right)^2 \quad (19)$$

From the above three independent equations (17)-(19) five unknowns parameters  $A, B, \rho, w, \gamma$  can be seen. For a better and deterministic solution two more condition is essential.

First, assuming that the scalar expansion ( $\theta$ ) is proportional to the shear scalar ( $\sigma$ ) and using the condition (16) gives

$$\frac{1}{\sqrt{3}}\left(\frac{\dot{A}}{A} - \frac{\dot{B}}{B}\right) = \alpha_0 \left(\frac{\dot{A}}{A} + 2\frac{\dot{B}}{B}\right) \quad (20)$$

where  $\alpha_0$  is a constant and the above equation yields to

$$\frac{\dot{A}}{A} = l\frac{\dot{B}}{B} \quad (21)$$

where  $l = \frac{2\alpha_0\sqrt{3}+1}{1-\alpha_0\sqrt{3}}$ . From the equation (21) gives

$$A = c_3 B^l \quad (22)$$

where,  $c_3$  is a constant and without the loss of generality taking  $c_3 = 1$ , yields

$$A = B^l \quad (23)$$

Secondly, a gauge function is considered as

$$\phi = \phi_0 a^\alpha = \phi_0 V^{\frac{\alpha}{3}} \quad (24)$$

A characterized scale factor is assumed as

$$a(t) = c_2 e^{\frac{t}{c_1}} (c_1 t + c_1 - 1)^{\frac{1}{c_1^2}} \quad (25)$$

where  $c_1$  and  $c_2$  are constants.

Using this equation (25) in (15), the deceleration parameter is obtained as

$$q = -1 + \frac{1}{(t+1)^2} \quad (26)$$

Using (16), (23) and (25) the metric functions are obtained as

$$A = k_3 e^{\frac{3lt}{c_1(l+2)}} (c_1 t + c_1 - 1)^{\frac{3l}{c_1^2(l+2)}} \quad (27)$$

$$B = k_1 e^{\frac{3t}{c_1(l+2)}} (c_1 t + c_1 - 1)^{\frac{3}{c_1^2(l+2)}} \quad (28)$$

$$C = k_2 e^{\frac{3t}{c_1(l+2)}} (c_1 t + c_1 - 1)^{\frac{3}{c_1^2(l+2)}} \quad (29)$$

where,  $k_1 = m^{\frac{-1}{l+2}} c_2^{\frac{3}{(l+2)}}$ ,  $k_2 = mk_1$ ,  $k_3 = k_1^l$ .

Hence, the metric (1) becomes

$$ds^2 = -dt^2 + k_3^2 e^{\frac{6lt}{c_1(l+2)}} (c_1 t + c_1 - 1)^{\frac{6l}{c_1^2(l+2)}} dx^2 + k_1^2 e^{\frac{6t}{c_1(l+2)}} (c_1 t + c_1 - 1)^{\frac{6}{c_1^2(l+2)}} e^{2x} dy^2 + k_2^2 e^{\frac{6t}{c_1(l+2)}} (c_1 t + c_1 - 1)^{\frac{6}{c_1^2(l+2)}} e^{-2x} dz^2 \quad (30)$$

#### 4. Physical and Geometrical Properties of the Model

The spatial volume ( $V$ ), Hubble's parameter ( $H$ ), expansion scalar ( $\theta$ ), shear scalar ( $\sigma$ ) and Anisotropic parameter ( $A_m$ ) are obtained as

$$V = c_2^3 e^{\frac{3t}{c_1}} (c_1 t + c_1 - 1)^{\frac{3}{c_1^2}} \quad (31)$$

$$H = \frac{1}{c_1} + \frac{1}{c_1} \frac{1}{(c_1 t + c_1 - 1)} \quad (32)$$

$$\phi = \phi_0 c_2^\alpha e^{\frac{\alpha t}{c_1}} (c_1 t + c_1 - 1)^{\frac{\alpha}{c_1^2}} \quad (33)$$

$$\theta = \frac{3(t+1)}{(c_1t+c_1-1)} \tag{34}$$

$$\sigma^2 = 3 \left(\frac{l-1}{l+2}\right)^2 \left(\frac{t+1}{c_1t+c_1-1}\right)^2 \tag{35}$$

$$A_m = 2 \left(\frac{l-1}{l+2}\right)^2 \tag{36}$$

From eqs. (19), (27), (28), and (33) the energy density, the EoS parameter and the skewness parameter are obtained as

$$\rho = \phi_0^2 c_2^{2\alpha} e^{\frac{2\alpha t}{c_1}} (c_1t + c_1 - 1)^{\frac{2\alpha}{c_1^2}} \left[ \frac{9(2l+1)(t+1)^2}{(l+2)^2(c_1t+c_1-1)^2} - k_4 \frac{1}{e^{c_1(l+2)}} \frac{1}{(c_1t+c_1-1)c_1^2(l+2)} + \frac{\alpha\omega}{2} \left( \frac{1}{c_1} + \frac{1}{c_1^2(c_1t+c_1-1)} \right)^2 \right] \tag{37}$$

where  $k_4 = \frac{1}{k_1^2 l}$

$$w = - \frac{\frac{27(t+2)^2-6}{(l+2)^2(c_1t+c_1-1)^2} + k_4 \frac{1}{e^{c_1(l+2)}} \frac{1}{(c_1t+c_1-1)c_1^2(l+2)} + \frac{\alpha\omega}{2} \left( \frac{1}{c_1} + \frac{1}{c_1^2(c_1t+c_1-1)} \right)^2}{\frac{9(2l+1)(t+1)^2}{(l+2)^2(c_1t+c_1-1)^2} - k_4 \frac{1}{e^{c_1(l+2)}} \frac{1}{(c_1t+c_1-1)c_1^2(l+2)} + \frac{\alpha\omega}{2} \left( \frac{1}{c_1} + \frac{1}{c_1^2(c_1t+c_1-1)} \right)^2} \tag{38}$$

$$\gamma = \frac{\frac{9(t+1)^2(l^2+l+1)}{(l+2)^2(c_1t+c_1-1)^2} + \frac{3-3l(c_1t+c_1-1)}{(l+2)(c_1t+c_1-1)^2} - k_4 \frac{1}{e^{c_1(l+2)}} \frac{1}{(c_1t+c_1-1)c_1^2(l+2)} + \frac{\alpha\omega}{2} \left( \frac{1}{c_1} + \frac{1}{c_1^2(c_1t+c_1-1)} \right)^2}{\frac{9(2l+1)(t+1)^2}{(l+2)^2(c_1t+c_1-1)^2} - k_4 \frac{1}{e^{c_1(l+2)}} \frac{1}{(c_1t+c_1-1)c_1^2(l+2)} + \frac{\alpha\omega}{2} \left( \frac{1}{c_1} + \frac{1}{c_1^2(c_1t+c_1-1)} \right)^2} \tag{39}$$

### 5. The Jerk parameter (j) and Statefinder Parameters {r, s}

The jerk parameter (j) is a dimensionless and third derivative of the scale factor with respect to cosmic time t [29,30]. It is defined as

$$j(t) = \frac{\ddot{a}}{aH^3} \tag{40}$$

Equation (40) can be written as

$$j(t) = q + 2q^2 - \frac{q}{H} \tag{41}$$

From equation (26), (32) and (41), yields

$$j(t) = 1 - \frac{3}{(t+1)^2} + \frac{2c_1}{(t+1)^2} \tag{42}$$

For flat  $\Lambda$ CDM model, jerk parameters (j) has the value  $j = 1$ . Sahni *et al.* [30] introduced the statefinder {r, s}, defined as

$$r = \frac{\ddot{a}}{aH^3} = 1 + 3 \frac{\dot{H}}{H^2} + \frac{\ddot{H}}{H^3} \tag{43}$$

$$\text{and } s = \frac{r-1}{3\left(q-\frac{1}{2}\right)} \tag{44}$$

From the above eqs. (26), (32), (43) and (44) the results obtained are

$$r = 1 - \frac{3}{(t+1)^2} + \frac{2c_1}{(t+1)^3} \tag{45}$$

$$\text{and } s = \frac{-12(t+1)-4c_1}{3(t+1)\{2-3(t+1)^2\}} \quad (46)$$

From the above results we see that as  $t \rightarrow \infty$ ,  $\{r, s\} \rightarrow \{1, 0\}$  which draw the idea that the model of the universe starts from Einstein static era to the  $\Lambda$ CDM model as in recent observation by Ahmad and Pradhan [31].

## 6. Conclusion

In this paper, a Bianchi Type-VI<sub>0</sub> cosmological model with a special form of scale factor in Sen-Dunn theory has been studied and solved for exact solutions. Spatial volume increases as time tends to infinity. Gauge function  $\varphi$  is also increasing function of time and increases as time passes. The Hubble parameter decreases and approaches to zero as time tend to infinity and our models anisotropic and homogeneous since  $\frac{\sigma^2}{\theta^2} = \text{constant}$ . Also the EoS parameter  $w$  is negative throughout the time  $t$  representing the DE model dynamics which in good agreement with recent observation. The jerk parameter ( $j$ ) has the value  $j = 1$ , deceleration parameter  $q = -1$ . A deceleration to acceleration transition occurs for models as time passes. This solution gives a scenario of the dark energy model which tends to a  $\Lambda$ CDM model.

## References

1. C. H. Brans and R. H. Dicke, Phys. Rev. **124**, 925 (1961).  
<https://doi.org/10.1103/PhysRev.124.925>
2. D. K. Sen and K. A. Dunn, J. Math. Phys. **12**, 578 (1971). <https://doi.org/10.1063/1.1665623>
3. K. A. Dunn, J. Math. Phys. **15**, 2229 (1974). <https://doi.org/10.1063/1.1666603>
4. D. R. K. Reddy, J. Phys. A: Mathematical Nucl. Gen. **6**, 12 (1973).  
<https://doi.org/10.1088/0305-4470/6/12/010>
5. D. R. K. Reddy, J. Math. Phys. **20**, 23 (1979). <https://doi.org/10.1063/1.523957>
6. A. R. Roy and B. Chatterjee, Ind. J. Pure Appl. Math. **12**, 211 (1981).
7. T. Singh and L. N. Rai, Gen. Relat. Gravit. **15**, 875 (1983).  
<https://doi.org/10.1007/BF00778798>
8. B. Mukherjee, Acta Phys. Hung. **A21/1**, 39 (2004). <https://doi.org/10.1556/APH.21.2004.1.4>
9. D. R. K. Reddy and R. L. Naidu, Astrophys. Space Sci. **312**, 99 (2007).  
<https://doi.org/10.1007/s10509-007-9657-7>
10. R. Venkateswarlu, J. Satish, and K. P. Kumar, The Afr. Rev. Phys. **8**, 169 (2013).
11. R. Venkateswarlu, J. Satish, and K. P. Kumar, Presapcetime J. **4**, 387 (2013).
12. H. R. Ghate and A. S. Sontakke, Global J. Inc. (USA), **14** (2014).
13. R. N. Patra, IJMTT **48**, 152 (2017). <https://doi.org/10.1002/piuz.201770315>
14. S. R. Roy and J. P. Singh, Acta Physica Austriaca **55**, 57 (1983).
15. L. K. Patel and S. S. Koppa, J. Aust. Math. Soc. Series B **33**, 77 (1991).  
<https://doi.org/10.1017/S033427000008638>
16. R. Bali, P. Pradhan, and A. Hassan, Int. J. Theo. Phys. **47**, 2594 (2008).  
<https://doi.org/10.1007/s10773-008-9694-1>
17. R. Bali, R. Banerjee, and S. K. Banerjee, Astrophys. Space Sci. **317**, 21 (2008).  
<https://doi.org/10.1007/s10509-008-9848-x>
18. A. Pradhan and R. Bali, Electron. J. Theo. Phys. **48**, 1466 (2008).  
<https://doi.org/10.1007/s10773-008-9918-4>
19. A. Asgar and M. Ansari, J. Theo. Appl. Phys. **8**, 219 (2014). <https://doi.org/10.1007/s40094-014-0151-7>



20. H. Amirhashchi, A. Pradhan, and D. S. Chouhan, *EJTP*. **11**, 109 (2014).
21. B. Mishra and P. K. Sahoo, *Astrophys. Space Sci.* **349**, 491 (2014).  
<https://doi.org/10.1007/s10509-013-1652-6>
22. M. V. Santhi, V. U. M.Rao, and Y. Aditya, *Can. J. Phys.* **95**, 179 (2017).  
<https://doi.org/10.1139/cjp-2016-0628>
23. M. V. Santhi, Y. Sobhanbabuand B. J. M. Rao, *J. Phys.: Conf. Ser.* **1344**, ID 012038 (2019).  
<https://doi.org/10.1088/1742-6596/1344/1/012038>
24. E. A. Hegazy, *Iran. J. Sci. Technol., Trans. A: Sci.* **43**, 663 (2019).  
<https://doi.org/10.1007/s40995-018-0530-z>
25. S. Satish and R. Venkateswarlu, *Bulg. J. Phys.* **46**, 67 (2019).
26. S. R. Bhojar, V. R. Chirde, and S. H. Shekh, *J. Sci. Res.* **11**, 249 (2019).  
<https://doi.org/10.3329/jsr.v11i3.39220>
27. M. Dewri, *J. Sci. Res.* **12**, 251 (2020). <https://doi.org/10.3329/jsr.v12i3.43313>
28. B. P. Brahma and M. Dewri, *High Technol. Lett.* **26**, 1049 (2020).  
<https://doi.org/10.37896/HTL26.08/1614>
29. T. Chiba and T. Nakamura, *Prog. Theor. Phys.* **100**, 1077 (1998).  
<https://doi.org/10.1143/PTP.100.1077>
30. V. Sahni, T. D. Saini, A. A. Starobinsky, and U. Alam, *J. Exp. Theor. Phys. Lett.* **77**, 201 (2003). <https://doi.org/10.1134/1.1574831>
31. N. Ahmad and A. Pradhan, *Int. J. Theo. Phys.* **53**, 289 (2014). <https://doi.org/10.1007/s10773-013-1809-7>

## ORIGINAL ARTICLE

 OPEN ACCESS

Received: 09-05-2022

Accepted: 31-07-2022

Published: 07-09-2022

**Citation:** Basumatary D, Dewri M (2022) Magnetized Bianchi Model with Time-dependent Deceleration Parameter. Indian Journal of Science and Technology 15(35): 1703-1711. <https://doi.org/10.17485/IJST/v15i35.1000>

\* **Corresponding author.**

[derphung17@gmail.com](mailto:derphung17@gmail.com)

**Funding:** None

**Competing Interests:** None

**Copyright:** © 2022 Basumatary & Dewri. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Published By Indian Society for Education and Environment (iSee)

**ISSN**

Print: 0974-6846

Electronic: 0974-5645

# Magnetized Bianchi Model with Time-dependent Deceleration Parameter

Derphungshar Basumatary<sup>1\*</sup>, Mukunda Dewri<sup>1</sup>

<sup>1</sup> Department of Mathematical Sciences, Bodoland University, Kokrajhar, 783370, BTR, Assam, India

## Abstract

**Objectives:** To investigate the Bianchi Type-III cosmological model in the presence of a magnetic field using the time-dependent deceleration parameter in Sen-Dunn scalar-tensor theory. **Methods:** In this work, we approach a Bianchi type-III metric with the presence of a magnetic field in the energy-momentum tensor. The time-dependent deceleration parameter representing an accelerated expansion proposed by Banerjee and Das (Gen. Relativ. Gravit. 37:10, 2005) is applied to obtain the exact solution of the field equations. Also, to obtain a deterministic solution considering shear scalar proportional to scalar expansion resulting in  $B = C^\beta$ . Furthermore, taking the power law relation between the scalar field and the average scale factor, the model's following physical and geometrical properties are computed and discussed with the present observational data. **Findings:** The model is anisotropic, expanding, shearing, and non-rotating with increasing volume as time tends to infinity from zero volume at  $t = 0$ . The model approves an accelerating universe under the  $\Lambda$ CDM model. **Novelty:** The study provides a better comprehension of cosmic evolution within the Sen-Dunn scalar-tensor gravitational theory context. The model demonstrates its consistency with the cosmological observation.

**Keywords:** Bianchi type III metric; Electromagnetic field; Deceleration parameter; SenDunn theory;  $\Lambda$ CDM model

## 1 Introduction

The most prominent study made by supernovae (SNe), cosmic microwave background (CMB), and baryonic acoustic oscillations (BAO) in modern cosmology is the accelerated expansion of the universe. Still, the universe holds the greatest mysteries of its evolution, such as the accelerated expansion's origin and cause. Many cosmologists take many modified theories that alternate with Einstein's theory into account to understand the accelerated expansion of the universe. Researchers are paying attention to these alternative theories because it is thought to offer a natural gravitational alternative to dark energy, which explains the universe's dark energy and late-time cosmic acceleration. Dark energy is highly considered the main element for accelerated expansion. Many studies have been executed for dark energy models in different gravitational theories in both the isotropic and anisotropic backgrounds. Cosmologists

employ isotropy as the most common and important type of spacetime to investigate how our universe (particularly DE) is currently behaving. However, the anisotropic models help us better understand the early stages of the universe's evolution. The finding of some anisotropy in the background radiation has now increased the importance of spatially homogenous and anisotropic Bianchi models. The two fluids, barotropic fluid and dark energy filled in Bianchi type-III universe were explored by Zia et al. <sup>(1)</sup> while considering the scale factor that results in a variable deceleration parameter. The EoS parameter emerges as the possible characteristic to study the nature of dark energy. In both interacting and non-interacting forms of the fluid, the equation of state for dark energy represents the phantom model at late times. For the non-interacting one, it begins in the quintessence region and moves through the  $\Lambda$ CDM model and eventually to the phantom model. Srivastava et al. <sup>(2)</sup> examined the anisotropic Bianchi type-III model with time-dependent deceleration parameter in general relativity; here, the EoS parameter for the new holographic dark energy (NHDE) falls under the k-essence region, which at late time coincides with the flat  $\Lambda$ CDM model. The Tsallis holographic dark energy model with time-dependent deceleration parameter in FRW universe addressed by Dixit et al. <sup>(3)</sup> effectively maintains the universe's acceleration. The EoS parameter, which corresponds to the suggested new holographic dark energy model, illustrates the change from quintessence to phantom. Further, the concept is also consistent with the flat  $\Lambda$ CDM model. In the Saez-Ballester modified theory of gravitation, Santhi and Sobhanbabu <sup>(4)</sup> also looked into the Tsallis holographic dark energy in Bianchi type III spacetime which coincides with the isotropic model discussed by Dixit et al. <sup>(3)</sup>. In the discussion of the anisotropic Bianchi Type-III dark energy cosmological model with a massive scalar meson field in general relativity, Raju et al. <sup>(5)</sup> noted that the EoS parameter of the model varies initially in the phantom region and then approaches the quintessence region as the model's physical parameters become independent of the scalar field at late times. In the case of modified holographic Ricci dark energy, Dixit et al. <sup>(6)</sup> explored the axially symmetric, spatially homogeneous anisotropic Bianchi type in Brans Dicke theory and noted that the EoS passes the phantom divide line and decreases with the increase in the scalar field. Many other authors investigated Bianchi type-III spacetime in different gravitational theories that provided many series of literature (Pawar and Sahare <sup>(7)</sup>, Koronur <sup>(8)</sup>, Bhardwaj and Rana <sup>(9)</sup>, Baro et al. <sup>(10)</sup>).

In cosmology, the magnetic field plays a crucial role in characterizing the ionized behavior that conducts energy fluctuations during the universe's expansion. The existence of the magnetic field affects the expansion independent of its strength and may result in anisotropy in the accelerated expansion (Matravers and Tsagas <sup>(11)</sup>). Furthermore, the impact of the magnetic field is slightly more significant in the early epoch; its influence cannot be excluded at a later cosmic phase around the present epoch (Ray et al. <sup>(12)</sup>). Even though the skewness disappears in the absence of a magnetic field, it does not do so when a magnetic field is present, although the magnetic field diminishes along with time (Sharma et al. <sup>(13)</sup>). Thus, the cosmos is expanding faster due to the decrease in magnetic permeability (Hegazy and Rehman <sup>(14)</sup>).

The Bianchi type-III spacetime is studied abundantly because of the unique geometric characteristics that set it apart from the other Bianchi types. The present work is motivated by the studies on Bianchi type-III spacetime in various contexts. In this study, we try to firm the knowledge of the Bianchi universe, considering the scalar-tensor theory of gravitation in the presence of an electromagnetic field. In order to obtain the exact solutions of the field equation, the special form of time-dependent deceleration parameter generalized by Banerjee and Das <sup>(15)</sup> is imposed to derive the model. The exact solution of the model is obtained considering the shear scalar proportional to the expansion scalar and the power law relation of the scalar field with the average scale factor. In this study, the model for the Bianchi type-III spacetime with the magnetic field in the scalar-tensor theory is also accelerating with exponential expansion approaching the standard  $\Lambda$ CDM model. In section 2, we formulate the metric and derive the field equations with the presence of an electromagnetic field. In section 3, the work presents the solutions to the field equation by considering some plausible conditions. In sections 4 and 5, the model's cosmological parameters are presented. In sections 6 and 7, we interpret and provide the study's conclusion.

## 2 Metric and Field Equations

The metric for Bianchi type-III considered as

$$ds^2 = -dt^2 + A^2 dx^2 + B^2 e^{-2mx} dy^2 + C^2 dz^2$$

where A, B, and C are the function of cosmic time t only.

The field equation given by Sen and Dunn <sup>(16)</sup> with gauge function (in natural units  $c = 1, 8\pi G = 1$ ) is

$$R_{ij} - \frac{1}{2} R g_{ij} = \omega \varphi^{-2} \left( \varphi_{,i} \varphi_{,j} - \frac{1}{2} g_{ij} \varphi_{,k} \varphi^{,k} \right) - \varphi^{-2} T_{ij}$$

where  $\omega = 3/2$ ,  $R_{ij}$  is the Ricci tensor, R is the Ricci scalar;  $T_{ij}$  is the energy-momentum tensor.

The energy-momentum tensor with the presence of an electromagnetic field is

$$T_{ij} = (\rho + p) u_i u_j + p g_{ij} + E_{ij}$$

Here  $u^i = (0, 0, 0, 1)$  is the four-velocity vector such that  $g^{ij} u_i u_j = -1$ . And taking

$$E_{ij} = \frac{1}{4\pi} [g^{hl} F_{ih} F_{jl} - \frac{1}{4} F_{hl} F^{hl} g_{ij}]$$

From the Maxwell equation,

$$\frac{\partial}{\partial x^j} (F^{ij} \sqrt{-g}) = 0$$

Eq. (5) leads to the result

$$F_{12} = K e^{-2mx}$$

where K and m are constants.

The field equation (2) with the equation (3), (4), and (6) for the equation (1) are

$$\frac{\ddot{A}}{A} + \frac{\ddot{B}}{B} + \frac{\dot{A}\dot{B}}{AB} - \frac{m^2}{A^2} = \frac{\omega}{2} \left(\frac{\dot{\phi}}{\phi}\right)^2 - \varphi^{-2} \left(p - \frac{K^2}{8\pi A^2 B^2}\right)$$

$$\frac{\ddot{B}}{B} + \frac{\ddot{C}}{C} + \frac{\dot{B}\dot{C}}{BC} = \frac{\omega}{2} \left(\frac{\dot{\phi}}{\phi}\right)^2 + \varphi^{-2} \left(p - \frac{K^2}{8\pi A^2 B^2}\right)$$

$$\frac{\ddot{A}}{A} + \frac{\ddot{C}}{C} + \frac{\dot{A}\dot{C}}{AC} = \frac{\omega}{2} \left(\frac{\dot{\phi}}{\phi}\right)^2 + \varphi^{-2} \left(p - \frac{K^2}{8\pi A^2 B^2}\right)$$

$$\frac{\ddot{A}}{A} + \frac{\dot{A}\dot{B}}{AB} + \frac{\dot{B}\dot{C}}{BC} - \frac{m^2}{A^2} = \varphi^{-2} \left(p + \frac{K^2}{8\pi A^2 B^2}\right) - \frac{\omega}{2} \left(\frac{\dot{\phi}}{\phi}\right)^2$$

$$m \left(\frac{\dot{A}}{A} - \frac{\dot{B}}{B}\right) = 0$$

The average scale factor for Bianchi type-III spacetime is

$$a(t) = (ABC)^{\frac{1}{3}}$$

The spatial volume, Hubble parameter is defined as

$$V = ABC = R^3$$

$$H = \frac{\dot{a}}{a} = \frac{1}{3} \left(\frac{\dot{A}}{A} + \frac{\dot{B}}{B} + \frac{\dot{C}}{C}\right)$$

And deceleration parameter is defined as

$$q = -\frac{a\ddot{a}}{\dot{a}^2}$$

The expansion scalar, shear scalar, and the anisotropic parameter are

$$\theta = 3H$$

$$\sigma^2 = \frac{1}{2} \sigma_{ij} \sigma^{ij} = \frac{1}{2} \left( \sum_{v=1}^3 H_v^2 - \frac{1}{3} \theta^2 \right)$$

$$A_m = \frac{1}{3} \sum_{v=1}^3 \left( \frac{H_v - H}{H} \right)^2$$

where,  $H_v(v = x, y, z)$  the directional parameter, and for the metric (1), it is given as  $H_x = \frac{\dot{A}}{A}$ ,  $H_y = \frac{\dot{B}}{B}$ ,  $H_z = \frac{\dot{C}}{C}$ .

### 3 Solutions of the field equations

From equation (11), by considering  $m = 1$ , we have

$$A = nB$$

Where  $n$  is the integration constant. Here we consider  $n = 1$  without the loss of generality. So the field equation (7)-(10) is transformed to

$$2 \frac{\ddot{B}}{B} + \left( \frac{\dot{B}}{B} \right)^2 - \frac{m^2}{B^2} = \frac{\omega}{2} \left( \frac{\dot{\phi}}{\phi} \right)^2 - \phi^{-2} \left( p - \frac{K^2}{8\pi A^2 B^2} \right)$$

$$\frac{\ddot{B}}{B} + \frac{\ddot{C}}{C} + \frac{\dot{B}\dot{C}}{BC} = \frac{\omega}{2} \left( \frac{\dot{\phi}}{\phi} \right)^2 + \phi^{-2} \left( p - \frac{K^2}{8\pi A^2 B^2} \right)$$

$$\frac{\ddot{B}}{B} + 2 \frac{\dot{B}\dot{C}}{BC} - \frac{m^2}{B^2} = \phi^{-2} \left( p + \frac{K^2}{8\pi A^2 B^2} \right) - \frac{\omega}{2} \left( \frac{\dot{\phi}}{\phi} \right)^2$$

Here, the nonlinear equation (20), (21), and (22) contains the unknown variables  $B, C, p, \rho$  and  $\phi$ . So to obtain the exact solution of the field equations above, we consider some plausible conditions.

To obtain the model compatible with the cosmological observation, we consider the variable deceleration parameter  $q$  (Banerjee and Das<sup>(15)</sup>) given by

$$q = -1 + \frac{k}{1 + a^k}$$

where " $a$ " is the scale factor, and  $k (> 0)$  is constant. The importance of the deceleration parameter lies in the hypothesis that the present universe has a transitional phase of expansion. It was decelerating in the past and is accelerating at present. Many researchers generalized the deceleration parameter to study the dynamical behavior of the accelerating universe. Pradhan et al.<sup>(17)</sup> utilized the time-dependent deceleration parameter certifying the accelerating universe and observing the Type-III model of the singularity of the Bianchi type-I universe in the background of  $f(R, T)$  gravity. The concept of bilinear varying deceleration parameter (BVDP), which shows super-exponential expansion, is highly discussed by Mishra and Chand<sup>(18)</sup> to extract the solutions of field equation for the anisotropic Bianchi type-I universe in the Saez-Ballester theory of gravitation. Recently, a deceleration parameter which is a function of time as well as periodic known as periodic time-varying deceleration parameter (PTVDP), which oscillates periodically from 2 to -4, representing accelerating expansion, has been used to observe the cosmological properties of the early universe as well as the fate of the universe by (Alam and Singh<sup>(19)</sup>, Garg et al.<sup>(20)</sup>).

From equation (23), we obtain the Hubble parameter as

$$H = \frac{\alpha e^{\alpha kt}}{e^{\alpha kt} - 1} = \alpha(1 + a^{-k})$$

where  $\alpha$  is an integrating constant.

As Collins et al. (21) expressed that for the spatially homogeneous metric, the normal congruence to the homogeneous hypersurface satisfies the condition  $\frac{\sigma}{\theta} = \text{constant}$ . This condition leads to

$$B = C^\beta$$

here,  $\beta$  is an arbitrary constant.

We consider the gauge function for the model as

$$\varphi = \varphi_0 \alpha^f(t)$$

where,  $\varphi_0$ ; and  $f$  being an ordinary constant.

From the equation (24) by integration, we have the scale factor as

$$a(t) = \left( e^{\alpha kt} - 1 \right)^{\frac{1}{k}}$$

Here,  $\alpha$  and  $k$  are positive constants.

Thus using equation (27) in equation (12), we get the following result as

$$A = B = \left( e^{\alpha kt} - 1 \right)^{\frac{3\beta}{k(2\beta+1)}}$$

$$C = \left( e^{\alpha kt} - 1 \right)^{\frac{3}{k(2\beta+1)}}$$

The model (1) with the equation (28) and (29) reduces to

$$ds^2 = -dt^2 + \left( e^{\alpha kt} - 1 \right)^{\frac{3\beta}{k(2\beta+1)}} dx^2 + \left( e^{\alpha kt} - 1 \right)^{\frac{3\beta}{k(2\beta+1)}} e^{-2mx} dy^2 + \left( e^{\alpha kt} - 1 \right)^{\frac{3}{k(2\beta+1)}} dz^2$$

#### 4 Cosmological Parameters of the model

From the equation (20)-(22), we obtain the following energy density and pressure using equations (26), (28), and (29)

$$\rho = \varphi_0^2 \left( e^{\alpha kt} - 1 \right)^{\frac{f}{k}} \left[ \frac{9(1+3\beta-\beta^2)\alpha^2 e^{2\alpha kt}}{(2\beta+1)^2 (e^{\alpha kt} - 1)^2} - \frac{3k\alpha^2 e^{\alpha kt}}{(2\beta+1)(e^{\alpha kt} - 1)^2} + \frac{3\alpha^2 f^2 e^{2\alpha kt}}{4(e^{\alpha kt} - 1)^2} \right] + \frac{K^2}{(e^{\alpha kt} - 1)^{\frac{12\beta}{k(2\beta+1)}}}$$

$$-p = \frac{\varphi_0^2}{2} \left( e^{\alpha kt} - 1 \right)^{\frac{f}{k}} \left[ \frac{9(1+3\beta+4\beta^2)\alpha^2 e^{2\alpha kt}}{(2\beta+1)^2 (e^{\alpha kt} - 1)^2} - \frac{3(3\beta+1)k\alpha^2 e^{\alpha kt}}{(2\beta+1)(e^{\alpha kt} - 1)^2} - m^2 \left( e^{\alpha kt} - 1 \right)^{\frac{-6\beta}{(2\beta+1)k}} - \frac{3\alpha^2 f^2 e^{2\alpha kt}}{2(e^{\alpha kt} - 1)^2} \right]$$

The spatial volume, expansion scalar, shear scalar, and anisotropy parameter for the model are as follows

$$V = \left( e^{\alpha kt} - 1 \right)^{\frac{3}{k}}$$

$$\theta = \frac{3\alpha e^{\alpha kt}}{e^{\alpha kt} - 1}$$

$$\sigma = \frac{\sqrt{3}(\beta-1)\alpha e^{\alpha kt}}{(2\beta+1)(e^{\alpha kt} - 1)}$$

$$A_m = \frac{2(\beta - 1)^2}{(2\beta + 1)^2} \tag{36}$$

Thus we also have

$$\frac{\sigma^2}{\theta^2} = \frac{(\beta - 1)^2}{3(2\beta + 1)^2} = \text{constant} (\neq 0, \text{ for } \beta > 1) \tag{37}$$

Energy condition and Statefinder parameter

$$\rho + p = \frac{\varphi_0^2}{2} (e^{\alpha kt} - 1)^{\frac{f}{k}} \left[ \frac{9(1 + 5\beta - 6\beta^2) \alpha^2 e^{2\alpha kt}}{(2\beta + 1)^2 (e^{\alpha kt} - 1)^2} + \frac{3(3\beta - 1)k\alpha^2 e^{\alpha kt}}{(2\beta + 1)(e^{\alpha kt} - 1)^2} + m^2 (e^{\alpha kt} - 1)^{\frac{-6\beta}{(2\beta + 1)k}} + \frac{3\alpha^2 f^2 e^{2\alpha kt}}{2(e^{\alpha kt} - 1)^2} \right] + \frac{K^2}{(e^{\alpha kt} - 1)^{\frac{12\beta}{k(2\beta + 1)}}} \tag{38}$$

$$\rho - p = \frac{\varphi_0^2}{2} (e^{\alpha kt} - 1)^{\frac{f}{k}} \left[ \frac{9(1 + 4\beta + 3\beta^2) \alpha^2 e^{2\alpha kt}}{(2\beta + 1)^2 (e^{\alpha kt} - 1)^2} + \frac{9(\beta - 1)k\alpha^2 e^{\alpha kt}}{(2\beta + 1)(e^{\alpha kt} - 1)^2} - m^2 (e^{\alpha kt} - 1)^{\frac{-6\beta}{(2\beta + 1)k}} \right] + \frac{K^2}{(e^{\alpha kt} - 1)^{\frac{12\beta}{k(2\beta + 1)}}}$$

$$\rho + 3p = \frac{\varphi_0^2}{2} (e^{\alpha kt} - 1)^{\frac{f}{k}} \left[ \frac{9(3\beta - 14\beta^2 - 1) \alpha^2 e^{2\alpha kt}}{(2\beta + 1)^2 (e^{\alpha kt} - 1)^2} + \frac{3(9\beta + 1)k\alpha^2 e^{\alpha kt}}{(2\beta + 1)(e^{\alpha kt} - 1)^2} + m^2 (e^{\alpha kt} - 1)^{\frac{-6\beta}{(2\beta + 1)k}} + \frac{6\alpha^2 f^2 e^{2\alpha kt}}{(e^{\alpha kt} - 1)^2} \right] + \frac{K^2}{(e^{\alpha kt} - 1)^{\frac{12\beta}{k(2\beta + 1)}}} \tag{40}$$

From the following result, the energy condition identified that the Null Energy Condition (NEC), the Weak Energy Condition (WEC), and the Strong Energy Condition (SEC) are satisfied throughout the cosmic evolution of the model. The model's Dominant Energy Condition (DEC) rapidly increases from negative phase to positive boundary after some cosmic time  $t = 0.3$ , and later it is satisfied for the stability of the universe's expansion.

The statefinder parameter  $\{r,s\}$  introduced by Shani et al. (22) is given as

$$r = \frac{\ddot{a}}{aH^3} = 1 + 3\frac{\dot{H}}{H^2} + \frac{\ddot{H}}{H^3}$$

$$s = \frac{(r - 1)}{3(q - \frac{1}{2})}$$

Here, H is the Hubble parameter, and q is the deceleration parameter. The cosmological diagnostic pair  $\{r,s\}$  allow us to determine the characteristic properties of the dark energy in a model-independent approach. With the equations (23) and (24), we can rewrite the diagnostic pair  $\{r,s\}$  as

$$r = 1 - \frac{3k}{e^{\alpha kt}} + \frac{k^2 (e^{\alpha kt}) + 1}{e^{2\alpha kt}}$$

$$s = \frac{2k (ke^{\alpha kt} + k - 3e^{\alpha kt})}{3e^{\alpha kt} (2k - 3e^{\alpha kt})}$$

We observe from the above result that when  $t \rightarrow \infty$  we get the pair  $(r, s) \rightarrow \{1, 0\}$ , which explains that the model of the universe starts from the radiation era to the  $\Lambda$ CDM model.

### 5 Results and Discussion

The equation (30) represents the model for the Bianchi type-III cosmological model with the electromagnetic field in the framework of Sen-Dunn scalar-tensor theory. The physical and geometrical parameters for the model are discussed above:




Fig 1. Volume Vs. Time: For  $\alpha = 0.5, k = 1.75$




Fig 2. Hubble parameter, Expansion scalar Vs. Time: For  $\alpha = 0.5, k = 1.75$

- The plotting of the figures are drawn against time (Gyr) by considering the values  $\alpha = 0.5, \beta = 1.5, k = 1.75, f = -0.62, \varphi_0 = 1$ . The graphical representation of the physical and kinematical parameters with this values signifies that the model obtained in this study are constrained with the recent cosmological observations.
- The model is expanding, shearing, and non-rotating, with the spatial volume increasing from the finite volume at  $t = 0$  and expanding to an infinite volume as  $t \rightarrow \infty$  is shown in Figure 1. The model’s expansion scalar and Hubble parameter positively decrease with the increase in time, as shown in Figure 2, which shows that the universe’s expansion rate is slower as time increases and expansion gradually ends for  $t \rightarrow \infty$ .




Fig 3. Deceleration Parameter Vs. Time

- The relation  $\frac{\sigma^2}{\theta^2} = constant$  show that the model (30) does not tend to isotropy for  $\beta \neq 1$  at late times. Its contrast to the Bianchi type I spacetime model considering a similar deceleration parameter proclaims isotropy for large values of "t" by Tiwari and Beesham (23).
- The model’s energy density and the pressure given by equation (31) and (32) positively decreases. Also, it tends to be zero as  $t \rightarrow \infty$ ; for a significant value of time  $t$ , as shown in Figure 4. A similar characteristic of the energy density, pressure, and Hubble parameter and expansion scalar is also observed in the string cosmological model for Bianchi type-III spacetime in Lyra geometry (Baro et al. (10)). The  $q$  bounds to the value  $-1 \leq q < 0$  from decelerating phase  $q > 0$  to de-Sitter expansion






Fig 4. Energy density, Pressure Vs. Time: For  $\alpha = 0.5, k = 1.75, \beta = 1.5, f = -0.62, \varphi_0 = 1$

or exponential expansion, which explains the transition from the early deceleration to the late time acceleration as shown in Figure 3 . The result of the deceleration parameter follows the recent cosmological data. Some model also exhibits super-exponential expansion at the early phase with  $q < -1$  and, finally, at late times, tends to exponential expansion  $q = -1$  (Santhi and Sobanbabu (4)).




Fig 5. Energy condition Vs. Time:For  $\alpha = 0.5, k = 1.75, \beta = 1.5, f = -0.62, \varphi_0 = 1$

- Initially, the dominant energy condition (DEC) gets violated for time  $t < 0.3$ , and then rapidly, it satisfies the energy condition as positively decreasing with time. The NEC, WEC, and SEC satisfy the energy condition throughout the time  $t$  (Figure 5). The energy condition depicts the early deceleration to the present acceleration of the universe.
- From the above result (43) and (44), the nature of the statefinder parameter is obtained as  $\{r, s\} \rightarrow (1, 0)$  as  $t \rightarrow \infty$ , which shows that the universe approaches the  $\Lambda$ CDM model at late times, which matches the recent cosmological model of the universe. The same result was observed in the previous study considering a special form of scale factor for Bianchi type-VI<sub>0</sub> in the framework of Sen-Dunn theory (Basumatary and Dewri (24)).

## 6 Conclusion

This paper investigates the solutions of Bianchi type-III spacetime in the presence of an electromagnetic field in the Sen-Dunn scalar-tensor theory of gravitation. To deliver the solutions to the field equations, we consider the time-dependent deceleration parameter  $q$ , a signature flip property (Banerjee and Das (15)). The model (30) expands with zero volume at  $t = 0$  to infinite as time  $t \rightarrow \infty$  and eventually, at late times, tends to the de-Sitter universe. For the significant value of  $t$ , the present model does not tend to isotropy, except for particular exceptional cases for  $\beta = 1$ . The model is expanding, shearing, non-rotating, and anisotropic throughout the evolution admitting initial singularity at  $t = 0$ , which shows a relevant characteristic of the model investigated by Pawar and Sahare (7). The diagnostic pair  $\{r, s\}$  represents that the model of the universe tends to  $\Lambda$ CDM as time  $t \rightarrow \infty$ , which is consistent with the recent observational data. The study presents a toy piece of the Bianchi type-III cosmological model in the Sen-Dunn scalar-tensor theory of gravitation. The generated and given model represents an expanding and accelerating cosmological model, delivered by examining the above physical and geometrical factors. The model studied in this part is compatible with the recent cosmological observation. Therefore, more profound knowledge of the cosmic evolution of the universe will result from studying the Bianchi spacetime in the scalar-tensor theory of gravitation.

## References

1) Zia R, Sharma UK, Maurya DC. Transit two-fluid models in anisotropic Bianchi type-III space-time. *New Astronomy*. 2019;72:83-91. Available from: <https://doi.org/10.1016/j.newast.2019.04.003>.

- 2) Srivastava S, Sharma UK, Pradhan A. New holographic dark energy in bianchi- III universe with k-essence. *New Astronomy*. 2019;68:57–64. Available from: <https://doi.org/10.1016/j.newast.2018.11.002>.
- 3) Dixit A, Sharma UK, Pradhan A. Tsallis holographic dark energy in FRW universe with time varying deceleration parameter. *New Astronomy*. 2019;73:101281–101281. Available from: <https://doi.org/10.1016/j.newast.2019.101281>.
- 4) Santhi MV, Sobhanbabu Y. Bianchi type-III Tsallis holographic dark energy model in Saez–Ballester theory of gravitation. *The European Physical Journal C*. 2020;80(12):1198–1198. Available from: <https://doi.org/10.1140/epjc/s10052-020-08743-9>.
- 5) Raju KD, Aditya Y, Rao VUM, Reddy DRK. Bianchi type-III dark energy cosmological model with massive scalar meson field. *Astrophysics and Space Science*. 2020;365(2):45–45. Available from: <https://doi.org/10.1007/s10509-020-03753-1>.
- 6) Dixit A, Singhal S, Zeyauddin M. Model for Modified Holographic Ricci Dark Energy in Gravitation Theory of Branc Dicke. *Walailak Journal of Science and Technology (WJST)*. 2021;18(3):6986–6986. Available from: <https://doi.org/10.48048/wjst.2021.6986>.
- 7) Pawar DD, Shahare SP. Dynamics of tilted Bianchi type-III cosmological model in  $f(R, T)$  gravity. *Journal of Astrophysics and Astronomy*. 2019;40(4):31–31. Available from: <https://doi.org/10.1007/s12036-019-9598-x>.
- 8) Korunur M. Tsallis holographic dark energy in Bianchi type-III spacetime with scalar fields. *Modern Physics Letters A*. 2019;34(37):1950310–1950310. Available from: <https://doi.org/10.1134/S020228932001003X>.
- 9) Bhardwaj VK, Rana MK. Bianchi–III Transitioning Space-Time in a Nonsingular Hybrid Universe within Lyra’s Cosmology. *Gravitation and Cosmology*. 2020;26(1):41–49. Available from: <https://doi.org/10.1134/S020228932001003X>.
- 10) Baro J, Singh KP, Singh AT. Mathematical analysis on anisotropic Bianchi type-III inflationary string cosmological models in Lyra geometry. *Indian Journal of Science and Technology*. 2021;14(1):46–54. Available from: <https://doi.org/10.17485/IJST/v14i1.1705>.
- 11) Matravers DR, Tsagas CG. Cosmic magnetism, curvature, and the expansion dynamics. *Physical Review D*. 2000;62(10):103519–103519. Available from: <https://doi.org/10.1103/PhysRevD.62.103519>.
- 12) Ray PP, Mishra B, Tripathy SK. Dynamics of anisotropic dark energy universe embedded in one-directional magnetized fluid. *International Journal of Modern Physics D*. 2019;28(07):1950093–1950093. Available from: <https://doi.org/10.1142/S0218271819500937>.
- 13) Sharma UK, Mishra AK, Pradhan A. A comparative study on Kaluza-Klein model with magnetic field in Lyra manifold and general relativity. *New Astronomy*. 2019;70:27–35. Available from: <https://doi.org/10.1016/j.newast.2019.02.003>.
- 14) Hegazy EA, Rahaman F. Bianchi type VI0 cosmological model with electromagnetic field and variable deceleration parameter in general theory of relativity. *Indian Journal of Physics*. 2020;94(11):1847–1852. Available from: <https://doi.org/10.1007/s12648-019-01614-4>.
- 15) Banerjee N, Das S. Acceleration of the universe with a simple trigonometric potential. *General Relativity and Gravitation*. 2005;37(10):1695–1703. Available from: <https://doi.org/10.1007/s10714-005-0152-6>.
- 16) Sen DK, Dunn KA. A Scalar-Tensor Theory of Gravitation in a Modified Riemannian Manifold. *Journal of Mathematical Physics*. 1971;12(4):578–586. Available from: <https://doi.org/10.1063/1.1665623>.
- 17) Pradhan A, Tiwari RK, Beesham A, Zia R. LRS Bianchi type-I cosmological models with accelerated expansion in  $f(R, T)$  gravity in the presence of  $\Lambda(T)$ . *The European Physical Journal Plus*. 2019;134(5):229–229. Available from: <https://doi.org/10.1140/epjp/i2019-12583-4>.
- 18) Mishra RK, Chand A. Cosmological models in Saez–Ballester theory with bilinear varying deceleration parameter. *Astrophysics and Space Science*. 2020;365(4):76–76. Available from: <https://doi.org/10.1007/s10509-020-03790-w>.
- 19) Alam MK, Singh SS. Interacting anisotropic dark energy with time dependent inhomogeneous equation of state. *International Journal of Nonlinear Analysis and Applications*. 2021;12:2167–2180. Available from: <https://doi.org/10.22075/IJNAA.2021.6076>.
- 20) Garg P, Pradhan A, Zeyauddin M. Roles of a periodic time varying deceleration parameter in particle creation with  $(d+2)$  dimensional FLRW Universe. *New Astronomy*. 2022;95:101821–101821. Available from: <https://doi.org/10.1016/j.newast.2022.101821>.
- 21) Collins CB, Glass EN, Wilkinson DA. Exact spatially homogeneous cosmologies. *General Relativity and Gravitation*. 1980;12(10):805–823. Available from: <https://dx.doi.org/10.1007/bf00763057>.
- 22) Sahni V, Saini TD, Starobinsky AA, Alam U. Statefinder—A new geometrical diagnostic of dark energy. *Journal of Experimental and Theoretical Physics Letters*. 2003;77(5):201–206. Available from: <https://doi.org/10.1134/1.1574831>.
- 23) Tiwari RK, Beesham A. Anisotropic model with decaying cosmological term. *Astrophysics and Space Science*. 2018;363(11):243–243. Available from: <https://doi.org/10.1007/s10509-018-3456-1>.
- 24) Basumatary D, Dewri M. Bianchi Type- $VI_{-5}$  Cosmological Model with Special Form of Scale Factor in Sen-Dunn Theory of Gravitation. *Journal of Scientific Research*. 2021;13(1):137–143. Available from: <https://doi.org/10.3329/jsr.v13i1.48479>.

**NATIONAL CONFERENCE ON  
EMERGING TRENDS IN ADVANCED MATHEMATICAL  
SCIENCES & ITS INTERDISCIPLINARY AREAS**

**21-22 February, 2020**



ORGANIZED BY

DEPARTMENT OF MATHEMATICAL SCIENCES  
BODOLAND UNIVERSITY, KOKRAJHAR, ASSAM, INDIA, PIN-783370



Certified that Dr/Mr/Ms... *Serphungshar Basumatary*.....

Department of... *Mathematical Sciences*.....

✓  
University/College/Institute... *Bodoland University, Kokrajhar*.....

has participated/presented a paper entitled... *Bianchi Type - IX Cosmological Model*.....

in... *Gen - Sum Theory of Gravitation*.....

in the "NATIONAL CONFERENCE ON EMERGING TRENDS IN ADVANCED MATHEMATICAL SCIENCES & ITS INTERDISCIPLINARY AREAS" held at Bodoland University, Kokrajhar, Assam, India.

Dr. Prahalad Basumatary  
Registrar  
Bodoland University

Prof. K. Priyokumar Singh  
(Convener)  
Dean, Faculty of Science & Technology  
Bodoland University

Prof. Laishram Ladusingh  
Vice-Chancellor  
Bodoland University



# CERTIFICATE

## NATIONAL CONFERENCE ON ADVANCED MATHEMATICAL & APPLIED SCIENCES

1<sup>st</sup> & 2<sup>nd</sup> July 2020



Organized by  
Department of Mathematical Sciences  
Bodoland University, Kokrajhar

*This is to certify that*

### **DERPHUNGSHAR BASUMATARY**

*a Research Scholar of Department of Mathematical Sciences, Bodoland University, Kokrajhar has presented a paper titled "Bianchi Type-VI<sub>0</sub> Cosmological Model with Special Form of Scale Factor in Sen-Dunn Theory of Gravitation" in the online mode National Conference on Advanced Mathematical & Applied Sciences held through Cisco Webex Meetings on 1<sup>st</sup> & 2<sup>nd</sup> July, 2020.*

**Dr. Rajshekhar Roy Baruah**  
Joint Convener

**Dr. Mukunda Dewri**  
Chairman & HoD i/c

**Prof. Laishram Ladusingh**  
Vice-Chancellor, Bodoland University

**Dr. Dipankar Das**  
Joint Convener



**NATIONAL CONFERENCE  
ON  
SCIENCE & TECHNOLOGY FOR SUSTAINABLE DEVELOPMENT  
(STSD-2022)**



Organized jointly by

**SCIENCE COLLEGE, KOKRAJHAR & VIJNANA BHARATI, NESM**

In collaboration with

**IASST, Guwahati, India & NECTAR, Shillong, India**



**Certificate of Presentation**

**This is to certify that**



*Mr. Derphungshar Basumatary*

of

**Bodoland University, Kokrajhar**

has presented a paper entitled "*Magnetized Bianchi Type-VI dark energy model with electromagnetic field*" in the National Conference on "Science & Technology for Sustainable Development (STSD-2022)" held on 9th-10th September, 2022 at Science College, Kokrajhar, Assam, India.

Principal  
Science College, Kokrajhar

President  
Vijnana Bharati, NESM

Chairperson  
National Conference  
STSD-2022

Convener  
National Conference  
STSD-2022

Convener  
National Conference  
STSD-2022

Organizing Secretary  
National Conference  
STSD-2022

