

Total No. of printed pages = 12

63/2 (SEM-3) PHY 303,304(N/O)

2021

(held in 2022)

PHYSICS

(Theory Paper)

Paper Code : PHY-303 (NEW)

(Advanced Nuclear Physics-I)

Full Marks – 80

Time – Two hours

The figures in the margin indicate full marks for the questions.

1. Choose the correct answer from the following :

1×5=5

(a) When a photon interacts with matter, pair production occurs if the energy E (in MeV) of the photon is

(i) $0.1 < E$

(ii) $0.1 < E < 10$

(iii) $E > 10$

(iv) None of the above

[Turn over

(b) In a typical range number-distance curve, the mean range is the thickness of the matter at which the number of particle absorbed is roughly

- (a) 10 % (b) 1%
(c) 50% (d) 100%

(c) Which of the following statement is correct ?

- (a) Λ is a baryon with positive charge.
(b) Intermediate vector bosons are composite in nature.
(c) e , μ and τ are known as mesons.
(d) Mesons and baryons are collectively known as hadrons.

(d) The range of electromagnetic force is

- (a) $\leq 10^{-14}$ cm (b) $\leq 10^{-13}$ cm
(c) $\sim 10^{-5}$ cm (d) infinite

(e) Choose the particle with strangeness quantum number, $s = 3$ from the following

- (i) $\bar{\Omega}^+$ (ii) Ξ^-
(iii) Ω^- (iv) Δ^+

2. Answer any *five* of the following : $2 \times 5 = 10$

- (a) Neutron is chargeless but it has a non-zero magnetic moment. — Why ?
(b) What is Schmidt line ? Write its significance.
(c) What are the different modes of nuclear shape vibration ? What is the lowest possible mode of shape vibration ?
(d) Define Detector sensitivity. Mention on which factors does Detector sensitivity depend.
(e) What are Absolute efficiency and Intrinsic efficiency ?
(f) What are the role of magnetic field and electric field in a cyclotron ?
(g) Explain the consequences when heavy ions are used in cyclotron.

3. Answer any *five* of the following : $5 \times 5 = 25$

- (a) Discuss the spin-parity of excited states of Te-120 nucleus in the light of vibrational model.

- (b) Why parity in rotational model is assumed to be positive? The first excited state of the rotational spectrum of the nucleus $^{238}_{92}\text{U}$ has an energy 45 KeV above the ground state. Calculate the energy (in KeV) of the second and third excited state.

$$1+4=5$$

- (c) Why the lowest order of electric multipole moment that may arise from a non-spherical charge distribution is the electric quadrupole moment. With the help of single particle shell model calculate the quadrupole moment of $^{27}_{13}\text{Al}$.

- (d) Write down the Bethe-Bloch equation and describe each term in details. Mention limitations of Bethe-Bloch formula. $4+1=5$

- (e) What do you mean by dead time of a detector? Discuss about the models of dead time detector. Why the concept of dead time is important in the context of particle detector.

$$1+3+1=5$$

- (f) Write short note on Surface Barrier Detector.

5

- (g) A fixed frequency cyclotron has an oscillation frequency of 12 MHz and a dee radius of 0.55 meter. It is used to accelerate deuterium, $m_d = 3.34245 \times 10^{-27}$ kg, and $e = 1.6 \times 10^{-19}$ C. Calculate

(i) the Magnetic flux density

(ii) Energy to which deuterium is accelerated.

4. Answer any *four* of the following: $10 \times 4 = 40$

- (a) Discuss how nuclear magnetic moment is calculated with the help of single particle shell model. Using single particle shell model calculate the nuclear magnetic moment of $^{13}_6\text{C}$ and $^{15}_7\text{N}$ and compare it with the experimental value: $\mu(^{13}_6\text{C})_{\text{exp}} \sim 0.7\mu_N$; $\mu(^{15}_7\text{N})_{\text{exp}} \sim -0.28\mu_N$. Discuss Schmidt's approach to calculate the magnetic moment of a nucleus by taking the example of $^{13}_6\text{C}$ and $^{15}_7\text{N}$.

$$2+4+4=10$$

- (b) What are the different mechanisms by which a charged particle interacts when it passes through matter. Discuss the working of a scintillation counter and mention its advantages over other detectors. $4+6=10$

(c) Write in detail the principle, working, and construction for Van de Graaff generator. Which type of particles can be accelerated using Van de Graaff generator? 8+2=10

(d) Write in detail the principle, working and construction for the Betatron. How is the problem of "loss of resonance" resolved? 8+2=10

(e) (i) A baryon X decays by strong interaction as $X \rightarrow \Sigma^+ + \pi^- + \pi^0$, where Σ^+ is a member of the isospin triplet (Σ^+ , Σ^0 , Σ^-). What will be the third component of the isospin (I_3) of X? 3

(ii) A particle is a composite state of three quarks u, d, and s. What will be the electric charge, spin and strangeness of the particle? 3

(iii) Build up the structure of spin-1 mesons and spin 3/2 baryons in terms of quarks. 2+2=4

(f) (i) What is $\theta - \tau$ puzzle? Discuss how Lee and Yang solved the mystery. 2+2=4

(ii) Describe C. S. Wu's experiment which lead to the discovery of parity violation in weak interaction. 6

(Theory Paper)

Paper Code : PHY-304 (OLD)

(Nuclear Physics-II)

Full Marks - 80

Time - Two hours

The figures in the margin indicate full marks for the questions.

1. Choose the correct answer from the following : 1×5=5

(a) When a photon interacts with matter, pair production occurs if the energy E (in MeV) of the photon is

(a) $0.1 < E$ (b) $0.1 < E < 10$

(c) $E > 10$ (d) None of the above

(b) In elastic scattering the incident particle after striking the target-nucleus leaves with

(i) less energy (ii) greater energy

(iii) same energy (iv) zero energy

(c) In nuclear reaction, the physical quantity, which is *not* conserved is

- (i) total energy.
- (ii) angular momentum.
- (iii) charge.
- (iv) electric quadrupole moment.

(d) Which of the following statement is correct ?

- (a) Λ is a baryon with positive charge.
- (b) Intermediate vector bosons are composite in nature.
- (c) e , μ , and τ are known as mesons.
- (d) Mesons and baryons are collectively known as hadrons.
- (e) Choose the particle with strangeness quantum number, $s = 3$ from the following:
 - (i) $\bar{\Omega}^+$ (ii) E^-
 - (iii) Ω^- (iv) Δ^+

2. Answer any *five* of the following : $2 \times 5 = 10$

- (a) Neutron is chargeless but it has a non-zero magnetic moment. — Why?
- (b) What is Schmidt line? Write its significance.
- (c) What are the different modes of nuclear shape vibration? What is the lowest possible mode of shape vibration?
- (d) What are Absolute efficiency and Intrinsic efficiency?
- (e) What are Stripping and Pick-up reactions? Give one example of each.
- (f) What is the necessity of nuclear collective models?

3. Answer any *five* of the following questions :

$5 \times 5 = 25$

- (a) Write the success and failure of single particle shell model.
- (b) Discuss the spin-parity of excited states of $Te-120$ nucleus in the light of vibrational model.

- (c) Why parity in rotational model is assumed to be positive? The first excited state of the rotational spectrum of the nucleus ${}^{238}_{92}\text{U}$ has an energy 45 KeV above the ground state. Calculate the energy (in KeV) of the second and third excited state.

$$1+4=5$$

- (d) Why the lowest order of electric multipole moment that may arise from a non-spherical charge distribution is the electric quadrupole moment? With the help of single particle shell model calculate the quadrupole moment of ${}^{27}_{13}\text{Al}$.

- (e) Write down the Bethe-Bloch equation and describe each term in details. Mention limitations of Bethe-Bloch formula.

$$4+1=5$$

- (f) What do you mean by dead time of a detector? Discuss about the models of dead time detector. Why the concept of dead time is important in the context of particle detector?

$$1+3+1=5$$

- (g) Write short note on Surface Barrier Detector.

5

4. Answer any four of the following: $10 \times 4 = 40$

- (a) Discuss how nuclear magnetic moment is calculated with the help of single particle shell model. Using single particle shell model calculate the nuclear magnetic moment of ${}^{13}_6\text{C}$ and ${}^{15}_7\text{N}$ and compare it with the experimental value: $\mu({}^{13}_6\text{C})_{\text{exp}} \sim 0.7 \mu_N$; $\mu({}^{15}_7\text{N})_{\text{exp}} \sim -0.28 \mu_N$. Discuss Schmidt's approach to calculate the magnetic moment of a nucleus by taking the example of ${}^{13}_6\text{C}$ and ${}^{15}_7\text{N}$.

$$2+4+4=10$$

- (b) What are the different mechanisms by which a charged particle interacts when it passes through matter. Discuss the working of a scintillation counter and mention its advantages over other detectors.

$$4+6=10$$

- (c) (i) A baryon X decays by strong interaction as $X \rightarrow \Sigma^+ + \pi^- + \pi^0$, where Σ^+ is a member of the isospin triplet (Σ^+ , Σ^0 , Σ^-). What will be the third component of the isospin (I_3) of X?

3

- (ii) A particle is a composite state of three quarks u , d , and s . What will be the electric charge, spin and strangeness of the particle ? 3
- (iii) Build up the structure of spin-1 mesons and spin $3/2$ baryons in terms of quarks. $2+2=4$
- (d) (i) What do you mean by nuclear reaction ? Explain briefly different types of nuclear reactions with example. $1+5=6$
- (ii) Discuss the mechanism of formation of compound nucleus. 4
- (e) Write short notes on any *two* : $5+5=10$
- (i) G.M counter
- (ii) Multi-wire proportional chambers
- (iii) Multi-Channel Analyzer (MCA).